

# INVERSE PROBLEMS AND INDEX FORMULAE FOR DIRAC OPERATORS

YAROSLAV KURYLEV AND MATTI LASSAS

**Abstract:** *We consider a Dirac-type operator  $D_P$  on a vector bundle  $V$  over a compact Riemannian manifold  $(M, g)$  with a nonempty boundary. The operator  $D_P$  is specified by a boundary condition  $P(u|_{\partial M}) = 0$  where  $P$  is a projector which may be a non-local, i.e. a pseudodifferential operator. We assume the existence of a chirality operator which decomposes  $L^2(M, V)$  into two orthogonal subspaces  $X_+ \oplus X_-$ . Under certain conditions, the operator  $D_P$  restricted to  $X_+$  and  $X_-$  defines a pair of Fredholm operators which maps  $X_+ \rightarrow X_-$  and  $X_- \rightarrow X_+$  correspondingly, giving rise to a superstructure on  $V$ . In this paper we consider the questions of determining the index of  $D_P$  and the reconstruction of  $(M, g)$ ,  $V$  and  $D_P$  from the boundary data on  $\partial M$ . The data used is either the Cauchy data, i.e. the restrictions to  $\partial M \times \mathbb{R}_+$  of the solutions to the hyperbolic Dirac equation, or the boundary spectral data, i.e. the set of the eigenvalues and the boundary values of the eigenfunctions of  $D_P$ . We obtain formulae for the index and prove uniqueness results for the inverse boundary value problems. We apply the obtained results to the classical Dirac-type operator in  $M \times \mathbb{C}^4$ ,  $M \subset \mathbb{R}^3$ .*

**AMS classification:** 35J25, 58J45.

**Keywords:** Inverse boundary value problem, Dirac equation, boundary control, focusing sequences, Aharonov-Bohm effect, index theorem.

## 1. INTRODUCTION

Recent study of inverse problems has successfully shown that external measurements can be used to uniquely determine coefficients of various partial differential equations modeling macroscopic and microscopic phenomena. Examples of these are the paradigm problem, the inverse problem for the conductivity equation encountered in the impedance tomography, the inverse problem for the wave equation with an inhomogeneous wave speed, the inverse scattering problem for the Schrödinger operator (for different approaches see e.g. [3, 5, 6, 25, 31, 44, 45, 52]). In these problems the structure of the space is *a priori* known before measurements, being usually a known domain of an Euclidean space. Recently, many inverse problems have been generalized to the cases where the underlying space is not *a priori* known but is assumed to be some unknown smooth Riemannian manifold. This was followed by the change of the attitude from proving the uniqueness in the considered

inverse problem to describing the corresponding groups of transformations of the unknown object which preserve the measured data, e.g. isometries of Riemannian manifolds, gauge equivalences of differential operators, etc. From this point of view any anisotropic inverse problem, even in a given domain  $\Omega \subset \mathbb{R}^n$ , falls into this second category having a non-trivial group of transformations which consists of some boundary preserving diffeomorphisms of the domain, see e.g. [4, 7, 32, 35, 39, 49] and, for a detailed description of this approach, [29].

A natural task is to generalize this approach further to inverse problems on general vector bundles as, in the modern physics, physical phenomena are often modeled by equations on bundles. Indeed, the vector bundles are encountered from the early development of the relativistic quantum mechanics to the modern quantum field theory. Thus, in this paper our aim is to bring the mathematical study of inverse problems closer to the modern physics and to study inverse problems for one of its basic equations, the Dirac equation on a general vector bundle. Assuming the existence of a chirality operator, we consider a Dirac equation on an unknown bundle and reconstruct the underlying manifold, the bundle structure on it and also the coefficients of the equation up to a natural group of transformations, namely, the isometries of the manifold and bundlemorphisms of the bundle. To this end we further advance the techniques developed for the inverse problems for Maxwell's equation [36, 37, 38].

For previous results in inverse problems for the Dirac equation in the 1-dimensional case, see e.g. [14, 20, 24] and in the 3-dimensional case, see [21, 26, 27, 56]. In the 3-dimensional case all these papers deal with perturbations of the canonical Dirac operator  $D_0$  in  $\mathbb{R}^3$  or  $\Omega \subset \mathbb{R}^3$ ,

$$(1) \quad D_0 u = i \sum_{k=1}^3 \alpha_k \partial_k u + mc^2 \alpha_0 u, \quad u \in L^2(\mathbb{R}^3; \mathbb{C}^4),$$

where  $\alpha_\nu, \nu = 0, \dots, 3$ , are the standard Dirac matrices. Thus the structure of the underlying space and Dirac bundle was known *a priori*.

**1.1. Formulation of the problem.** In this paper we consider two types of inverse boundary value problems for the Dirac-type operator,  $D$ , acting on sections of a complex vector bundle,  $V$ , over a compact, connected Riemannian  $n$ -manifold  $(M, g)$  with a non-empty boundary,  $\partial M$ . Roughly speaking (for a rigorous formulation of the problems see Subsection 2.4 below), the first inverse problem deals with waves, i.e. the solutions to the hyperbolic problem,

$$(2) \quad (i\partial_t + D)u(x, t) = 0 \quad \text{in } M \times \mathbb{R}_+, \quad u|_{t=0} = 0.$$

Then the data used is the set of the Cauchy data of these waves restricted to  $\partial M \times \mathbb{R}_+$ , i.e. the set  $C_0(D) = \{u|_{\partial M \times \mathbb{R}_+} : u \text{ satisfies (2)}\}$ . The corresponding inverse problem, called the *dynamic inverse problem*, is that of the unique determination of the manifold  $M$ , the bundle

$V$ , and the operator  $D$  when we are given the boundary  $\partial M$ , the restriction  $W$  of the bundle to  $\partial M$ ,  $W = V|_{\partial M}$ , and the *dynamic boundary data*  $C_0(D)$ .

The second inverse problem or, more rigorously, a family of inverse problems, deals with the self-adjoint Dirac-type operator,

$$(3) \quad D_P u = Du; \quad \mathcal{D}(D_P) = \{u \in H^1(M, V) : P(u|_{\partial M}) = 0\}.$$

Here  $P$  is an orthoprojector in  $L^2(\partial M, W)$ , given by a zero-order classical pseudo-differential operator on  $\partial M$  which, in general, is not a local operator. The corresponding data are the *boundary spectral data*,

$$(\lambda_k, \phi_k|_{\partial M}), \quad k = 1, 2, \dots,$$

where  $\lambda_k, \phi_k$  are the eigenvalues and normalized eigenfunctions of  $D_P$ .

Throughout the paper, we assume the existence of a chirality operator  $F : V \rightarrow V$ , acting fiber-wise, i.e.,  $F : \pi^{-1}(x) \rightarrow \pi^{-1}(x)$ , with  $\pi : V \rightarrow M$  being the projector onto the base  $M$ . The chirality operator satisfies the anti-commutation relation

$$(4) \quad D \circ F + F \circ D = 0$$

and makes it possible, under some conditions, to split  $D_P$  into a pair of operators  $D_P^\pm$ ,

$$D_P = \begin{pmatrix} 0 & D_P^- \\ D_P^+ & 0 \end{pmatrix}.$$

We will derive formulae to express  $\text{Ind}(D_P^\pm)$  in terms of the dynamical boundary data.

The main tool in our study of the inverse problems for the Dirac-type equations is the boundary control method, see [5], or, more precisely, its geometric version, see [29] for a detailed exposition. The method is based on the finite velocity of the wave propagation and control properties of the waves generated by boundary sources.

The plan of the paper is as follows: In Section 2 we give definitions of the Dirac bundles and Dirac-type operators, introduce a decomposition of  $L^2(M, V)$  into two channels  $X_+ \oplus X_-$  and formulate rigorously the main results of the paper. In Section 3 we describe properties of the hyperbolic Dirac-type equation (2) and provide a unique continuation result. In Section 4 we derive a formula for the energy of a wave in each channel in terms of the boundary data and prove some global and local boundary controllability results for the hyperbolic Dirac-type equation. Generalized boundary sources and an extension of the time-derivative  $\partial_t$  are introduced in Section 5. The relation of  $\partial_t$  and  $D_P$  is studied in Section 6 where we obtain index formulae for  $D_P$  in terms of the boundary data. In Section 7 we prove that the dynamic inverse data determine the manifold  $M$  and its metric  $g$ . In Section 8 we introduce focusing sources which generate waves that, at given time, are localized at a single point  $x \in M$ . These waves with different polarizations are used in the reconstruction of the bundle  $V$  and operator  $D_P$  from the

dynamic inverse data. In Section 9 we study the high-energy asymptotics of the solutions to prove the uniqueness for the inverse boundary spectral problem. In Section 10 we apply the obtained results to the Dirac-type equation (1) in  $\mathbb{R}^3$  leading to an analog of the Aharonov-Bohm effect. In Appendix we prove, using the construction of Brüning and Lesch, the self-adjointness of the Dirac-type operator with the local boundary condition.

## 2. DEFINITIONS AND MAIN RESULTS.

**2.1. Definitions.** Here we introduce some basic definitions, results and examples about Dirac bundles and operators on them following the terminology of [13, 16, 55].

Let  $(M, g)$  be a compact connected  $C^\infty$ -smooth Riemannian  $n$ -manifold with a non-empty boundary. Let  $V$  be a smooth complex vector bundle over  $M$  and denote the projection onto the base manifold by  $\pi : V \rightarrow M$ . Each fiber  $\pi^{-1}(x)$  is a complex  $d$ -dimensional vector space with a Hermitian inner product  $\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_x$ . We define  $|\phi|_x^2 = \langle \phi, \phi \rangle_x$  and denote the smooth sections of  $V$  by  $C^\infty(M, V)$ . Endomorphisms  $\text{End}(V)$  are fiber-preserving smooth maps,  $L : V \rightarrow V$ ,  $\pi L(\phi) = \pi \phi$  that are linear in each fiber. By  $\text{Clif}(M)$  we denote the Clifford bundle over  $M$ . This means that, at each  $x \in M$ , the fiber  $\text{Clif}_x(M)$  is an algebra generated by the vectors in  $T_x M$  with a product  $\cdot$  satisfying the relation

$$(5) \quad v \cdot w + w \cdot v = -2g(v, w) \quad \text{for } v, w \in T_x M \subset \text{Clif}_x(M).$$

We assume that there is a fiber-wise map  $\gamma : \text{Clif}(M) \rightarrow \text{End}(V)$  that provides a Clifford module structure for  $V$ , i.e.  $\gamma$  gives an action, for any  $x \in M$ , of the algebra  $\text{Clif}_x(M)$  on fibers  $\pi^{-1}(x)$  of  $V$  and satisfies

$$(6) \quad \langle \gamma(v)\lambda, \mu \rangle_x + \langle \lambda, \gamma(v)\mu \rangle_x = 0, \quad v \in T_x(M), \quad \lambda, \mu \in \pi^{-1}(x).$$

Let  $\nabla$  be the Levi-Civita connection in  $(M, g)$ . It defines a connection on  $\text{Clif}(M)$  that satisfies

$$\nabla_X(v \cdot w) = (\nabla_X v) \cdot w + v \cdot (\nabla_X w),$$

where  $X \in C^\infty(M, TM)$  and  $v, w \in C^\infty(M, \text{Clif}(M))$ . We denote by  $\nabla$  also a connection on  $V$  that is compatible with  $\langle \cdot, \cdot \rangle$  and  $\gamma$ , i.e.,

$$\begin{aligned} X \langle \phi, \psi \rangle &= \langle \nabla_X \phi, \psi \rangle + \langle \phi, \nabla_X \psi \rangle, \\ \nabla_X(\gamma(w)\phi) &= \gamma(\nabla_X w)\phi + \gamma(w)\nabla_X \phi, \end{aligned}$$

where  $X \in C^\infty(M, TM)$ ,  $w \in C^\infty(M, \text{Clif}(M))$ , and  $\phi, \psi \in C^\infty(M, V)$ . When these conditions are satisfied, we say that  $(V, \langle \cdot, \cdot \rangle, \gamma, \nabla)$  is a Dirac bundle over  $(M, g)$ , for brevity, a Dirac bundle  $V$ .

The unperturbed Dirac operator,  $D_0$  on the Dirac bundle  $V$ , which is sometimes called the *Dirac operator associated with a Clifford connection*, is locally given by

$$(7) \quad D_0 u|_U = \sum_{j=1}^n \gamma(e_j) \nabla_{e_j} u|_U,$$

where  $(e_1, \dots, e_n)$  is an orthonormal frame in an open set  $U \subset M$ .

As noted before, we assume that there is a chirality operator  $F : V \rightarrow V$  acting fiber-wise and satisfying

$$(8) \quad \begin{aligned} \langle F\lambda, \mu \rangle_x &= \langle \lambda, F\mu \rangle_x, \quad \lambda, \mu \in \pi^{-1}(x), \\ \gamma(v) \circ F + F \circ \gamma(v) &= 0, \quad v \in T_x M, \quad F^2 = I, \\ \nabla F &= 0, \quad \text{i.e.} \quad \nabla_X(Fu) = F(\nabla_X u) \end{aligned}$$

where  $X \in C^\infty(M, TM)$  and  $u \in C^\infty(M, V)$ . When  $n = \dim(M)$  is even, a chirality operator always exists, namely,

$$(9) \quad F\phi|_U = (\sqrt{-1})^{n/2} \gamma(e_1) \gamma(e_2) \dots \gamma(e_n) \phi|_U.$$

A chirality operator exists also when we deal with a space-type hypersurface on a Lorentzian manifold, e.g. [23].

Using a chirality operator, we define the fiber-wise orthogonal projectors onto the  $+1$  and  $-1$  eigenspaces of  $F$ ,

$$(10) \quad \Pi_+ = \frac{1}{2}(I + F), \quad \Pi_- = \frac{1}{2}(I - F).$$

We denote by  $Q \in \text{End}(V)$  a self-adjoint potential, i.e.,

$$(11) \quad \langle Q\lambda, \mu \rangle_x = \langle \lambda, Q\mu \rangle_x, \quad \lambda, \mu \in \pi^{-1}(x),$$

which respects the chirality structure, i.e.

$$(12) \quad F \circ Q + Q \circ F = 0.$$

We call the minimal Dirac-type operator,  $D_{\min}$ , the operator

$$(13) \quad D_{\min} u = D_0 u + Qu, \quad u \in \mathcal{D}(D_{\min}) = C_0^\infty(M, V),$$

where  $C_0^\infty(M, V)$  is the space of the smooth compactly supported section of  $V$ .

**Example 1.** As an example, we consider the so-called form-Dirac operator, see e.g. [16, 55]. Let  $(M, g)$  be an oriented Riemannian manifold of dimension  $n$  and  $\Omega^j M$  be the space of the complex-valued differential  $j$ -forms on  $M$ . Let  $\Omega M = \Omega^0 M \oplus \dots \oplus \Omega^n M$  be the Grassmannian bundle on  $M$  and  $*$  the Hodge operator of  $\Omega(M)$  related to the metric  $g$ . For  $\lambda \in \Omega_x^j M$  and  $\mu \in \Omega_x^k M$ , let  $\langle \lambda, \mu \rangle_x = 0$  if  $j \neq k$  and  $\langle \lambda, \mu \rangle_x = *(\lambda \wedge *\bar{\mu})$  for  $j = k$ .

Denote by  $I : T_x M \rightarrow T_x^* M$  the identification  $I(a^j \frac{\partial}{\partial x^j}) = g_{jk} a^j dx^k$  and by  $\iota_v : \Omega_x^j M \rightarrow \Omega_x^{j-1} M$  the inner product with a vector  $v \in T_x M$ , e.g.

$$\iota_v w(v_1, \dots, v_{j-1}) = w(v, v_1, \dots, v_{j-1}),$$

for any  $w \in \Omega_x^j M$ ,  $v, v_1, \dots, v_{j-1} \in T_x M$ . Then the map  $\gamma : T_x M \rightarrow \text{End}(\Omega_x M)$ ,

$$\gamma(v)\lambda = -I(v) \wedge \lambda + \iota_v \lambda$$

extends to a homomorphism  $\gamma : \text{Clif}_x(M) \rightarrow \text{End}(\Omega_x M)$ . Let  $\nabla$  be the Levi-Civita connection on  $\Omega M$ . Then  $(\Omega M, \langle \cdot, \cdot \rangle, \gamma, \nabla)$  is a Dirac bundle. The unperturbed Dirac operator (7) on this bundle is

$$D_0 : \Omega M \rightarrow \Omega M, \quad D_0 = d + \delta : \Omega^j M \rightarrow \Omega^{j+1} M \oplus \Omega^{j-1} M,$$

where  $d$  is the exterior differential and

$$\delta = (-1)^{nj+1} * d* : \Omega^j M \rightarrow \Omega^{j-1} M$$

is the codifferential. A chirality operator,  $F$  for the form-Dirac operator may be defined simply as

$$(14) \quad F\lambda = (-1)^j \lambda, \quad \lambda \in \Omega^j M,$$

splitting the Grassmannian bundle into the forms of the even and odd orders,  $\Omega M = \Omega^e M \oplus \Omega^o M$ .

**Example 2.** Consider Maxwell's equations in  $M \times \mathbb{R}$ ,  $M \subset \mathbb{R}^3$ ,

$$\begin{aligned} \text{curl } E(x, t) &= -B_t(x, t), & D(x, t) &= \epsilon(x)E(x, t), \\ \text{curl } H(x, t) &= D_t(x, t), & B(x, t) &= \mu(x)H(x, t) \end{aligned}$$

where  $\epsilon(x)$  and  $\mu(x)$  are positive definite matrix valued functions. The velocity of the wave propagation is independent of the wave polarization if and only if  $\mu(x) = \alpha(x)^2 \epsilon(x)$  with some scalar function  $\alpha$ . In this case the travel time is determined by the Riemannian metric  $g_{ij} = \alpha^{-2} \det(\epsilon)^{-1} \delta^{ik} \epsilon_k^j$ . As shown in [36, 38], Maxwell's system may be extended to the hyperbolic equation

$$(15) \quad i\partial_t \omega + i(d - \delta_\alpha)\omega = 0, \quad \omega(x, t) = (\omega^0, \omega^1, \omega^2, \omega^3) \in \Omega M,$$

where  $\delta_\alpha \omega^j = \alpha \delta(\alpha^{-1} \omega^j)$  and  $\delta$  is the codifferential with respect to the travel time metric  $g$ . Indeed, a solution of Maxwell's system gives rise to a solution of the Dirac-type equation (15) with  $\omega^0 = 0$ ,  $\omega^1 = E_j dx^j$ ,  $\omega^2 = B_3 dx^1 \wedge dx^2 - B_2 dx^1 \wedge dx^3 + B_1 dx^2 \wedge dx^3$ , and  $\omega^3 = 0$ . Defining the inner product

$$\langle \omega^j, \lambda^k \rangle_x = \alpha(x)^{-1} * (\omega^j \wedge * \bar{\lambda}^k) \delta^{jk}, \quad \text{for } \omega^j \in \Omega_x^j M, \lambda^k \in \Omega_x^k M,$$

the operator  $i(d - \delta_\alpha)$  is formally self-adjoint in  $\Omega M$ . This operator defines a Dirac-type bundle  $(\Omega M, \langle \cdot, \cdot \rangle_x, \tilde{\gamma}, \nabla)$  where  $\tilde{\gamma}(v)\lambda = -i(I(v) \wedge \lambda + \iota_v \lambda)$  and  $\nabla$  is the Levi-Civita connection in  $(M, g)$ . The chirality operator (14) is a chirality operator on this bundle, too.

**2.2. Self-adjoint operators.** We return now to the discussion of the boundary conditions used to extend the minimal Dirac operator (13) to a self-adjoint operator in  $L^2(M, V)$  with the inner product

$$\langle \langle \phi, \psi \rangle \rangle = \int_M \langle \phi(x), \psi(x) \rangle_x dV_g(x),$$

where  $dV_g$  is the Riemannian volume on  $(M, g)$ . Self-adjoint extensions of  $D_{\min}$  defined by *non-local* boundary conditions, go back to the centennial work by Atiyah-Patodi-Singer [1]. In this paper, we will work with the self-adjoint extensions of  $D_{\min}$  extensively studied by Brüning and Lesch [13], see also [16, 22].

In the following we denote by  $H^s(M, V)$  the Sobolev spaces of sections of  $V$  with components in the Sobolev spaces  $H^s(U)$  in local smooth trivializations  $\Phi_U : \pi^{-1}(U) \rightarrow U \times \mathbb{C}^d$ ,  $U \subset M$  and by  $H_0^s(M, V)$  the closure of  $C_0^\infty(M, V)$  in  $H^s(M, V)$ .

Using the natural embedding  $j : \partial M \rightarrow M$ , we introduce the induced bundle  $W = V|_{\partial M} = j^*V$  on  $\partial M$ .

Next we consider the possible extensions, in  $L^2(M, V)$ , of the minimal Dirac-type operator (13). We use  $D$  to denote the maximal extension of the operator  $D_{\min}$  to  $H^1(M, V)$ ,

$$(16) \quad Du = (D_0 + Q)u, \quad \mathcal{D}(D) = H^1(M, V).$$

It is shown in [13] that there is a wide class of self-adjoint extensions of  $D_{\min}$  defined by boundary conditions of form (3),

$$(17) \quad D_P u = Du, \quad \mathcal{D}(D_P) = \{u \in H^1(M, V) : P(u|_{\partial M}) = 0\},$$

so that  $D_P \subset D$ . Here  $P$  is a zero-order classical pseudo-differential operator defining an orthoprojector on  $L^2(\partial M, W)$ . It should satisfy the following conditions:

$$(18) \quad P\gamma(N) = \gamma(N)(I - P),$$

and

$$(19) \quad \{P, P_{APS}\} \text{ is a Fredholm pair,}$$

that is,  $P_{APS} : \text{Ran}(P) \rightarrow \text{Ran}(P_{APS})$  is a Fredholm operator. Here  $N$  stands for the unit interior normal field of  $\partial M$  and the operator  $P_{APS}$  is associated with the *Atiyah-Patodi-Singer* boundary condition. Namely, let  $A(0)$ , sometimes called the *hypersurface Dirac operator*, (see e.g. Appendix or [22]), be a self-adjoint operator in  $L^2(\partial M, W)$ ,

$$(20) \quad A(0) = -\gamma(N) \sum_{\alpha=1}^{n-1} \gamma(e_\alpha) \nabla_\alpha + \frac{n-1}{2} H(x'), \quad \nabla_\alpha = \nabla_{e_\alpha}.$$

Here the vector fields  $e_\alpha$ ,  $\alpha = 1, \dots, n-1$ , form a local orthonormal frame on  $\partial M$ , while  $H(x')$  is the mean curvature of  $\partial M$  at the point  $x'$ , where  $x' = (x_1, \dots, x_{n-1})$  are local coordinates on  $\partial M$ . Then  $P_{APS}$  is a spectral projector of  $A(0)$  such that

$$(21) \quad P_{(0,\infty)} \subset P_{APS} \subset P_{[0,\infty)},$$

$$(22) \quad P_{APS} = -\gamma(N)(I - P_{APS})\gamma(N),$$

where we denote by  $P_{(a,b)}$  the spectral projector of  $A(0)$  associated with an interval  $(a, b)$ .

**Theorem 2.1** (Brüning-Lesch). *Let  $P$  be a zero-order classical pseudo-differential operator defining an orthoprojector which satisfies conditions (18) and (19). Let  $D_P$  be the Dirac-type operator (17). Then  $D_P$  is self-adjoint, the spectrum of  $D_P$  is discrete, and all eigenspaces are finite dimensional. Moreover,  $D_P$  is regular, that is,  $D_P u \in H^s(M, V)$  implies that  $u \in H^{s+1}(M, V)$  for any  $s \geq 0$ .*

The proof of Theorem 2.1 is given in [13, Thm. 1.5] for more general Dirac-type operators than those considered in this paper and in [16, 22] for more restricted cases.

In the sequel we will denote the eigenvalues of  $D_P$ , numerated according to their multiplicity, by  $\lambda_k^P$ ,  $|\lambda_k^P| \leq |\lambda_{k+1}^P|$ , and the corresponding orthonormal eigenfunctions by  $\phi_k^P$ . When there is no danger of confusion we will skip the index  $P$  in these notations.

Using projections  $\Pi_+$  and  $\Pi_-$ , see (10) we define the bundles  $V_+ = \Pi_+ V$  and  $V_- = \Pi_- V$  over  $M$  with the projections to  $M$  denoted by  $\pi_\pm$ . Note that, for any  $x \in M$ ,  $\pi_+^{-1}(x)$  and  $\pi_-^{-1}(x)$  are orthogonal so that  $V = V_+ \oplus V_-$  and  $L^2(M, V) = L^2(M, \Pi_+ V) \oplus L^2(M, \Pi_- V)$ . With a slight abuse of notation we use  $\Pi_\pm$  for the orthoprojectors in  $L^2(M, V)$  onto  $L^2(M, \Pi_\pm V)$ . The maximal Dirac-type operator  $D$  satisfies  $DF + FD = 0$  so that  $D$  can be decomposed as

$$(23) \quad D : H^1(M, \Pi_+ V) \oplus H^1(M, \Pi_- V) \rightarrow L^2(M, \Pi_- V) \oplus L^2(M, \Pi_+ V),$$

$$D \begin{pmatrix} u_+ \\ u_- \end{pmatrix} = \begin{pmatrix} 0 & D \\ D & 0 \end{pmatrix} \begin{pmatrix} u_+ \\ u_- \end{pmatrix}, \quad u_\pm = \Pi_\pm u.$$

When

$$(24) \quad \Pi_+ \mathcal{D}(D_P) \subset \mathcal{D}(D_P) \quad \text{or, equivalently,} \quad \Pi_- \mathcal{D}(D_P) \subset \mathcal{D}(D_P),$$

decomposition (23) gives rise to a decomposition of  $D_P$ . Namely, for  $D_P^\pm = D_P : \Pi_\pm \mathcal{D}(D_P) \rightarrow L^2(M, \Pi_\mp V)$ ,

$$(25) \quad D_P : \Pi_+ \mathcal{D}(D_P) \oplus \Pi_- \mathcal{D}(D_P) \rightarrow L^2(M, \Pi_+ V) \oplus L^2(M, \Pi_- V),$$

$$D_P \begin{pmatrix} u_+ \\ u_- \end{pmatrix} = \begin{pmatrix} 0 & D_P^- \\ D_P^+ & 0 \end{pmatrix} \begin{pmatrix} u_+ \\ u_- \end{pmatrix}.$$

If (24) is valid and thus the decomposition (25) is possible, we say that the bundle  $V = V_+ \oplus V_-$  on  $M$  has a *superstructure*. It then follows from [13] that the resulting operators  $D_P^\pm$  are Fredholm with, in general, non-trivial indices. For important relations of the index and the geometry of the bundle, see e.g. [1, 9, 18, 19].

**2.3. Local boundary condition.** As noted in Introduction, the boundary projection  $P$  in (17) is usually non-local. However, for a Dirac-type operator with chirality, it is possible to introduce a *local boundary condition* which makes it self-adjoint. To this end, we consider a fiberwise operator  $\Gamma$  on  $W = V|_{\partial M}$ ,

$$(26) \quad \Gamma : W \rightarrow W, \quad \Gamma \lambda = F \gamma(N) \lambda, \quad \lambda \in \pi^{-1}(x), \quad x \in \partial M.$$



It is clear from properties (6) and (8) that

$$(27) \quad \begin{aligned} \Gamma^2 &= I, \quad \Gamma \circ F + F \circ \Gamma = 0, \\ \langle \Gamma \lambda, \mu \rangle_x &= \langle \lambda, \Gamma \mu \rangle_x, \quad \lambda, \mu \in \pi^{-1}(x). \end{aligned}$$

Let

$$(28) \quad P_\Gamma = \frac{1}{2}(I + \Gamma),$$

be an orthoprojector acting fiber-wise on  $\pi^{-1}(x)$ ,  $x \in \partial M$ . Then  $P_\Gamma$  satisfies conditions (19) and (18) and, therefore, defines a self-adjoint Dirac-type operator  $D_\Gamma := D_{P_\Gamma}$ . We note that this operator will play a crucial role in the sequel. Because of this, we give in the Appendix a sketch of the proof of its self-adjointness.

**Example 1, continued.** Clearly,  $P_\Gamma$  defines a local boundary condition for the form-Dirac operator  $d + \delta$  determined in Example 1. In addition, there are other local boundary condition closely related to the de Rham complexes over  $(M, g)$ . Recall that  $j : \partial M \rightarrow M$  is a natural embedding. The *relative boundary condition* is defined by

$$(29) \quad P_r(u|_{\partial M}) := j^*(u) = 0,$$

and the *absolute boundary condition* by

$$(30) \quad P_a(u|_{\partial M}) := j^*(\iota_N(u)) = 0.$$

In the future we denote the corresponding self-adjoint form-Dirac operators by  $(d + \delta)_r$  and  $(d + \delta)_a$ .

**Example 3.** Another important example of a Dirac operator on the Grassmannian bundle  $\Omega M$ , for  $n = 2k$ , is given by

$$D_1 u = (d + \delta)u, \quad u \in \mathcal{D}(D_1) = \{u \in H^1(M, V) : P_{APS}(u|_{\partial M}) = 0\},$$

where  $P_{APS}$  is the Atiyah-Patodi-Singer boundary condition defined in subsection 2.2, see e.g. [1, 19, 41]. This operator is called the signature operator. In this case the chirality operator is given by (9).

## 2.4. Boundary data for inverse problems.

**Definition 2.2.** Consider a Dirac bundle  $(V, \langle \cdot, \cdot \rangle, \gamma, \nabla)$  with chirality  $F$ . The induced bundle structure on  $\partial M$  is the collection

$$(31) \quad \{\partial M, g|_{\partial M}, W, \langle \cdot, \cdot \rangle_x|_{x \in \partial M}, \gamma(N)|_{\partial M}, F|_{\partial M}\}.$$

Here  $\partial M$  is considered as a Riemannian manifold with the differentiable structure induced by the embedding  $j : \partial M \rightarrow M$  and the metric  $g|_{\partial M} = j^*g$ . The bundle  $W = V|_{\partial M}$  is the induced bundle on  $\partial M$ ,  $\langle \cdot, \cdot \rangle_x$  the Hermitian structure to  $W$ ,  $\gamma(N)|_{\partial M}$  and  $F|_{\partial M}$  are the restrictions of the Clifford action  $\gamma(N)$  and of the chirality operator  $F$  on  $W$ .

**Definition 2.3.** Let  $D_P$  be a self-adjoint Dirac-type operator of form (17) on a Dirac bundle  $(V, \langle \cdot, \cdot \rangle, \gamma, \nabla)$  with chirality  $F$ . The set

$$(32) \quad \{(\lambda_k, \phi_k|_{\partial M})\}_{k=1}^\infty,$$

where  $\lambda_k$  and  $\phi_k$  are the eigenvalues and the orthonormal eigenfunctions of  $D_P$  is called the boundary spectral data of  $D_P$ .

To define another type of boundary data, observe that any Dirac-type operator (17) is associated with a hyperbolic initial-boundary value problem

$$(33) \quad (i\partial_t + D)u(x, t) = 0 \quad \text{in } M \times \mathbb{R}_+,$$

$$(34) \quad P(u|_{\partial M \times \mathbb{R}_+}) = f \in P\mathring{C}^\infty(\partial M \times \mathbb{R}_+, W), \quad u|_{t=0} = 0,$$

where  $\mathring{C}^\infty(\partial M \times \mathbb{R}_+, X)$  is the class of the smooth functions on  $\partial M \times \mathbb{R}_+$  (with values in  $X$ ) which are equal to 0 near  $t = 0$ . We denote by  $u = u^f$  the solution of (33)-(34). Problem (33)-(34) defines the response operator,  $\Lambda_P$ ,

$$(35) \quad \Lambda_P f = u^f|_{\partial M \times \mathbb{R}_+}.$$

When  $P = P_\Gamma$ , we denote  $\Lambda_\Gamma = \Lambda_{P_\Gamma}$ . Other data for the inverse boundary problem could be the response operator  $\Lambda_P$ . However, we prefer to work with data which is independent of  $P$ .

**Definition 2.4.** Let  $D$  be the maximal Dirac-type operator (16). Then the Cauchy data set of  $D$  is the set

$$(36) \quad C_0(D) = \{u|_{\partial M \times \mathbb{R}_+} : u \in \mathring{C}^\infty(M \times \mathbb{R}_+, V) \text{ satisfies (2)}\}.$$

Later we prove the following equivalence of different type of data.

**Lemma 2.5.** *Assume that the induced bundle structure (31) on  $\partial M$  is given. Then the Cauchy data set  $C_0(D)$  determines the map  $\Lambda_\Gamma$  and vice versa. Moreover, the set  $C_0(D)$  and a projector  $P$  satisfying (18) and (19) determine the map  $\Lambda_P$  and vice versa.*

**2.5. Main results.** First we formulate our main index formula.

**Theorem 2.6.** *Let  $V$  be a Dirac bundle over  $(M, g)$  with a chirality operator  $F$ . Let  $D_P$  be a self-adjoint Dirac-type operator of the form (17).*

*Assume that we are given the induced bundle structure (31) on  $\partial M$  and the response operator  $\Lambda_P$ . Then these data determine, in a constructive way, a Hilbert space  $\mathcal{F}$ , a pair of projectors  $B_+$  and  $B_-$  in  $\mathcal{F}$  with  $B_+ \oplus B_- = I$ , and an unbounded self-adjoint operator  $\partial_t : \mathcal{F} \rightarrow \mathcal{F}$  with  $\mathcal{D}(\partial_t) \subset \mathcal{F}$  such that*

1. *The condition (24) for  $D_P$  is valid if and only if  $B_+\mathcal{D}(\partial_t) \subset \mathcal{D}(\partial_t)$ .*
2. *If decomposition (24) is valid then the operator*

$$(37) \quad \partial_t^+ : B_+\mathcal{D}(\partial_t) \rightarrow B_-\mathcal{F}, \quad \partial_t^+ u = \partial_t u$$

is a Fredholm operator and

$$(38) \quad \text{Ind}(\partial_t^+) = \text{Ind}(D_P^+).$$

In Theorem 2.6 the Hilbert space  $\mathcal{F}$  is the completion of the functions  $PC_0^\infty(\partial M \times [0, T], W)$ , where  $T > 0$  is sufficiently large. This completion is taken with respect to a seminorm  $\|\cdot\|_{\mathcal{F}}$  that is explicitly determined by the Cauchy data set  $C_0(D)$  (see (50), (51), (73) below). The operator  $\partial_t$  in  $\mathcal{F}$  is an extension of the time derivative

$$\partial_t f = \frac{\partial f}{\partial t}, \quad \text{for } f \in PC_0^\infty(\partial M \times [0, T], W).$$

Next we formulate our main results on the reconstruction of the manifold and the Dirac bundle. To this end, let  $V$  and  $\tilde{V}$  be vector bundles over diffeomorphic manifolds  $M$  and  $\tilde{M}$ . Let  $\ell : M \rightarrow \tilde{M}$  be a diffeomorphism and  $\pi : V \rightarrow M$  and  $\tilde{\pi} : \tilde{V} \rightarrow \tilde{M}$  be projections on the base spaces. A map  $L : V \rightarrow \tilde{V}$  is called a *bundlemorphism* compatible with  $\ell$  if  $\ell(\pi\phi) = \tilde{\pi}(L\phi)$ ,  $\phi \in V$  and

$$L : \pi^{-1}(x) \rightarrow \tilde{\pi}^{-1}(\ell(x)) \text{ is a linear isomorphism for any } x \in M.$$

Dealing with Riemannian manifolds and Dirac bundles on them we need to extend the notation of a bundlemorphism.

**Definition 2.7.** Let  $(V, \langle \cdot, \cdot \rangle, \gamma, \nabla)$  and  $(\tilde{V}, \langle \cdot, \cdot \rangle, \tilde{\gamma}, \tilde{\nabla})$  be Dirac bundles over isometric Riemannian manifolds  $(M, g)$  and  $(\tilde{M}, \tilde{g})$  with chirality operators  $F$  and  $\tilde{F}$ , and let  $D$  and  $\tilde{D}$  be Dirac-type operators of form (16) on them. A bundlemorphism  $L$  compatible with an isometry  $\ell : M \rightarrow \tilde{M}$  is called a *Dirac bundlemorphism* if

$$(39) \quad \begin{aligned} \langle L\phi, L\psi \rangle_{\ell(x)} &= \langle \phi, \psi \rangle_x, \quad \phi, \psi \in \pi^{-1}(x), \\ \tilde{\gamma}(d\ell(v)) &= L\gamma(v)L^{-1}, \quad v \in T_x M, \\ \tilde{F} &= LFL^{-1}, \quad \tilde{D} = LDL^{-1}, \end{aligned}$$

where  $d\ell$  is the differential map of  $\ell$ .

We turn now to some induced structures on the boundary.

**Definition 2.8.** Let  $(V, \langle \cdot, \cdot \rangle, \gamma, \nabla)$  and  $(\tilde{V}, \langle \cdot, \cdot \rangle, \tilde{\gamma}, \tilde{\nabla})$  be Dirac bundles over manifolds  $M$  and  $\tilde{M}$  with  $W = V|_{\partial M}$  and  $\tilde{W} = \tilde{V}|_{\partial \tilde{M}}$ . Let  $\partial M$  and  $\partial \tilde{M}$  be isometric with an isometry  $\kappa : \partial M \rightarrow \partial \tilde{M}$ . We say that a bundlemorphism  $K : W \rightarrow \tilde{W}$  is an *isomorphism between the induced bundle structures* of  $\partial M$  and  $\partial \tilde{M}$  compatible with  $\kappa$  if

$$\begin{aligned} \langle K\phi, K\psi \rangle_{\kappa(x)} &= \langle \phi, \psi \rangle_x, \quad \tilde{F}_{\kappa(x)} = KF_x K^{-1}, \\ \tilde{\gamma}(d\kappa(v)) &= K\gamma(v)K^{-1}, \quad \tilde{\gamma}(\tilde{N}(\kappa(x))) = K\gamma(N(x))K^{-1}. \end{aligned}$$

Here  $x \in \partial M$ ,  $\kappa(x) \in \partial \tilde{M}$ ,  $\phi, \psi \in \pi^{-1}(x)$  and  $v \in T_x(\partial M)$ . At last,  $N(x)$  and  $\tilde{N}(\tilde{x})$  are the unit interior normal vectors to  $\partial M$  at  $x$  and to  $\partial \tilde{M}$  at  $\tilde{x}$ .

Let now  $u \in \mathring{C}^\infty(M \times \mathbb{R}_+, V)$  be a solution of the hyperbolic Dirac equation (33) and  $L : V \rightarrow \tilde{V}$  be a Dirac bundlemorphism. Then  $\tilde{u} = Lu$  is a solution of the hyperbolic Dirac equation (33) with  $D$  replaced with  $\tilde{D}$ . Therefore, the Cauchy data sets  $C_0(D)$ ,  $C_0(\tilde{D})$  satisfy

$$K(C_0(D)) = C_0(\tilde{D}),$$

with  $K = L|_W$ . This shows that the following uniqueness result is optimal.

**Theorem 2.9.** *Let  $D$  and  $\tilde{D}$  be Dirac-type operators of the form (16) on the Dirac bundles with chirality  $(V, F)$  and  $(\tilde{V}, \tilde{F})$ . Assume that there is an isometry  $\kappa : \partial M \rightarrow \partial \tilde{M}$  and an isomorphism between the induced bundle structures of  $\partial M$  and  $\partial \tilde{M}$ ,  $K : W \rightarrow \tilde{W}$ , which is compatible with  $\kappa$ . Assume also that the Cauchy data sets satisfy*

$$K(C_0(D)) = C_0(\tilde{D}).$$

*Then there is an isometry  $\ell : (M, g) \rightarrow (\tilde{M}, \tilde{g})$  with  $\ell|_{\partial M} = \kappa$  and a Dirac bundlemorphism  $L : V \rightarrow \tilde{V}$  with  $L|_W = K$ . In particular,  $\tilde{D} = LDL^{-1}$ .*

Note that this theorem states that the Dirac operators are equal up to a bundlemorphism, rather than the connections  $\nabla$  and  $\tilde{\nabla}$ .

For the inverse boundary spectral problem we obtain a similar result.

**Theorem 2.10.** *Let  $D_P$  and  $\tilde{D}_{\tilde{P}}$  be self-adjoint Dirac-type operators of form (17) on the Dirac bundles with chirality  $(V, F)$  and  $(\tilde{V}, \tilde{F})$ . Assume that there is an isometry  $\kappa : \partial M \rightarrow \partial \tilde{M}$ , an isomorphism  $K : W \rightarrow \tilde{W}$  compatible with  $\kappa$ , and that the eigenvalues and the normalized eigenfunctions of  $D_P$  and  $\tilde{D}_{\tilde{P}}$  satisfy*

$$\tilde{\lambda}_j = \lambda_j, \quad \tilde{\phi}_j \circ \kappa|_{\partial M} = K(\phi_j|_{\partial M}), \quad j \in \mathbb{Z}_+.$$

*Then there is an isometry  $\ell : (M, g) \rightarrow (\tilde{M}, \tilde{g})$  with  $\ell|_{\partial M} = \kappa$  and a Dirac bundlemorphism  $L : V \rightarrow \tilde{V}$  with  $L|_W = K$  such that*

$$\tilde{D}_{\tilde{P}} = LD_PL^{-1}, \quad \tilde{P} = KPL^{-1}.$$

**Remark 2.11.** To consider above results in more practical terms, assume that we are given the boundary  $\partial M$ , the bundle  $W$  on it, and the Cauchy data set  $C_0(D)$  or the boundary spectral data. The proofs of the above theorems are constructive, providing a method to recover the manifold  $M$  (up to an isometry), the bundle  $V$  with its Hermitian,  $\langle \cdot, \cdot \rangle$  and Clifford module,  $\gamma$  structures, and the operators  $D$  and  $F$  (up to a Dirac bundlemorphism).

### 3. HYPERBOLIC DIRAC EQUATION AND UNIQUE CONTINUATION

**3.1. Basic properties.** We start with a general boundary condition  $P$  satisfying (18) and (19). Denote by  $u^f(t) = u^f(x, t)$  the solution of the initial boundary value problem (33)-(34).

By the Bochner-Lichnerowicz formula, see e.g. [8, 11],

$$D_0^2 = \nabla^* \nabla + \mathcal{R},$$

where  $\mathcal{R}$  is the curvature endomorphism on  $V$ . Hence  $D^2 = \nabla^* \nabla + B(x, D)$ , where  $B(x, D)$  is a first order differential operator. Thus the hyperbolic Dirac-type equation in (33) implies the wave equation

$$(40) \quad (\partial_t^2 + \nabla^* \nabla + B(x, D))u^f(x, t) = 0.$$

In the rest of subsection 3.1 we consider the local boundary condition associated to operator  $P_\Gamma$ , cf. (28). Let  $u = u^f(x, t)$ ,  $t \in \mathbb{R}_+$  be the solution of

$$(41) \quad \begin{aligned} (i\partial_t + D)u(x, t) &= 0 \quad \text{in } M \times \mathbb{R}_+, \\ P_\Gamma(u|_{\partial M \times \mathbb{R}_+}) &= f, \quad u|_{t=0} = 0, \end{aligned}$$

where  $f \in \mathring{C}^\infty(\partial M \times \mathbb{R}_+, P_\Gamma W)$ .

Since  $D_\Gamma$  is a self-adjoint operator with  $\mathcal{D}(D_\Gamma) \subset H^1(M, V)$ , the map  $\mathcal{U} : f \mapsto u^f$  has an extension

$$\mathcal{U} : H_0^1(\partial M \times [0, T], P_\Gamma W) \rightarrow C([0, T], H^1(M, V)) \cap C^1([0, T], L^2(M, V)),$$

when  $T > 0$ .

Next we discuss the finite velocity of the wave propagation for the initial boundary value problem (41). To describe it, let  $\Sigma \subset \partial M$  be open. We define the *domain of influence* of  $\Sigma$  at time  $T$  as

$$M(\Sigma, T) = \{x \in M \mid \text{dist}(x, \Sigma) \leq T\},$$

where  $\text{dist}(x, y)$  is the distance in  $(M, g)$ . Using local trivializations of the Dirac bundle and applying classical results on the finite velocity of the wave propagation in hyperbolic systems due to e.g. Wilcox [58], we see that if  $\text{supp}(f) \subset \Sigma \times [0, T]$ , then

$$(42) \quad \text{supp}(u^f(T)) \subset M(\Sigma, T).$$

Finally, using the local boundary condition, we prove Lemma 2.5.

*Proof:* Assume that we are given  $C_0(D)$  and  $P$ . Then  $\Lambda_P$  is determined by its graph  $\{(Pu, u) : u \in C_0(D)\}$  and visa versa. Since the induced bundle structure on  $\partial M$  determines  $P_\Gamma$ ,  $C_0(D)$  determines  $\Lambda_\Gamma$ .  $\square$

**3.2. Unique continuation.** Our next goal is to supplement the finite velocity result by a unique continuation result of Tataru's type, see [53, 54] for the pioneering works. To this end, for any open  $\Sigma \subset \partial M$ , we define the double cone of influence,

$$K_{\Sigma, T} = \{(x, t) \in M \times \mathbb{R} : |t - T| + \text{dist}(x, \Sigma) < T\} \subset M \times \mathbb{R}.$$

**Theorem 3.1.** *Let  $\Sigma \subset \partial M$  be open,  $u \in C((0, 2T), L^2(M, V))$  be a weak solution of the hyperbolic equation  $(i\partial_t + D)u = 0$  in  $M \times (0, 2T)$ . Assume that*

$$(43) \quad u|_{\Sigma \times (0, 2T)} = 0.$$

*Then  $u(x, t) = 0$  for  $(x, t) \in K(\Sigma, T)$ .*

*Proof:* Assume first that  $u \in C^\infty(M \times (0, 2T), V)$ . Condition (43) and the fact that  $u$  solves (33) imply that

$$(44) \quad \partial_N u|_{\Sigma \times (0, 2T)} = 0,$$

where  $\partial_N$  is the normal derivative. By (40),  $u$  satisfies the wave equation which, in a local trivialisation  $\Phi : \pi^{-1}(U) \rightarrow U \times \mathbb{C}^d$  where  $U \subset M$  is open, may be written componentwise as

$$(45) \quad (\partial_t^2 + D^2)u = (\partial_t^2 - \Delta_g)Iu + \tilde{B}_1(x, \partial_x)u = 0.$$

Here  $\Delta_g$  is the scalar Laplace operator on  $(M, g)$ ,  $I$  is identity matrix, and  $\tilde{B}_1$  is a first-order operator. We intend to apply Tataru's unique continuation result [53], in order to infer from (43), (44), and (45) that  $u$  is zero in  $K(\Sigma, T)$ . The difficulty is that the proof in [53] deals only with a scalar wave equation. In the case of the scalar wave equation with time-independent coefficients a variant of the proof is given in [29, Section 2.5]. This proof can be extended, word by word, to the vector wave equations of the form  $(\partial_t^2 - \Delta_g)Iu + \tilde{B}_1u = 0$ , where the leading order term is a scalar operator times the identity matrix. Therefore, (43), (44), and (45) imply that  $u = 0$  in  $K(\Sigma, T)$ .

Let now  $u \in C((0, 2T), L^2(M, V))$  be a weak solution of the hyperbolic Dirac-type equation (33) with the boundary condition (43) understood in a weak sense. To prove the claim, we start by smoothing  $u$  with respect to time. Let  $\psi(s) \in C_0^\infty(-1, 1)$ ,  $\int \psi ds = 1$ . Then,

$$u_\varepsilon(x, t) = \int_{\mathbb{R}} u(x, t - s)\psi_\varepsilon(s)ds, \quad \psi_\varepsilon(s) = \frac{1}{\varepsilon}\psi\left(\frac{s}{\varepsilon}\right),$$

is in  $C^\infty((\varepsilon, 2T - \varepsilon), L^2(M, V))$  and solves there the Dirac-type wave equation  $(i\partial_t + D)u_\varepsilon = 0$  with  $u_\varepsilon|_{\Sigma \times (\varepsilon, 2T - \varepsilon)} = 0$ . As

$$D^2u_\varepsilon = -\partial_t^2u_\varepsilon \in C^\infty((\varepsilon, 2T - \varepsilon), L^2(M, V)),$$

the standard techniques of the elliptic regularity theory, e.g. [17], implies that  $u_\varepsilon \in C^\infty((\varepsilon, 2T - \varepsilon), H^1(M, V))$ . Iterating this analysis we see that  $u_\varepsilon \in C^\infty(M \times (\varepsilon, 2T - \varepsilon), V)$ . By the first part of the proof,

$$u_\varepsilon = 0 \quad \text{for } \{(x, t) : \text{dist}(x, \Sigma) + |t - T| < T - \varepsilon\}.$$

As  $u_\varepsilon \rightarrow u$ , when  $\varepsilon \rightarrow 0$ , this implies that  $u = 0$  in  $K(\Sigma, T)$ .  $\square$

## 4. INNER PRODUCTS AND CONTROLLABILITY.

**4.1. Inner products.** In this section we consider solutions  $u^f(t) = u^f(x, t)$  of (33)-(34),  $f \in P\mathring{C}^\infty(\partial M \times \mathbb{R}_+, W)$ . We compute the norms of  $u_+^f(t) = \Pi_+ u^f(t)$  and  $u_-^f(t) = \Pi_- u^f(t)$  using the boundary data, that is, the induced bundle structure (31) on  $\partial M$  and the operator  $\Lambda_P$ . This type of formulae are called the Blagovestchenskii formula, due to their first appearance in [10] for the one-dimensional inverse problems.

Using (8) and (12) we see that  $FD + DF = 0$  and thus  $\Pi_+ D = D\Pi_-$  and  $\Pi_- D = D\Pi_+$ . Thus, we can write the hyperbolic Dirac-type equation (33) as

$$(46) \quad \left( i\partial_t + \begin{pmatrix} 0 & D \\ D & 0 \end{pmatrix} \right) \begin{pmatrix} u_+(x, t) \\ u_-(x, t) \end{pmatrix} = 0 \quad \text{in } M \times \mathbb{R}_+,$$

where  $u_+(x, t) = \Pi_+ u(x, t)$  and  $u_-(x, t) = \Pi_- u(x, t)$ .

**Theorem 4.1.** *Let  $f, h \in P\mathring{C}^\infty(\partial M \times \mathbb{R}_+, W)$  be given and  $u^f, u^h$  be the solutions of (33)-(34). Then the response operator  $\Lambda_P$ , and the induced bundle structure (31) on  $\partial M$  determine the inner products*

$$(47) \quad \langle\langle u_+^f(t), u_+^h(s) \rangle\rangle \quad \text{and} \quad \langle\langle u_-^f(t), u_-^h(s) \rangle\rangle \quad \text{for } s, t \geq 0.$$

*In particular, these data determine  $\langle\langle u^f(t), u^h(s) \rangle\rangle$  for  $s, t \geq 0$ .*

*Proof:* We consider only the first inner product in (47) as the second one can be evaluated in the same manner. The Stokes' formula for the Dirac-type operator  $D$  takes the form,

$$(48) \quad \langle\langle D\phi, \psi \rangle\rangle - \langle\langle \phi, D\psi \rangle\rangle = \int_{\partial M} \langle \gamma(N)\phi, \psi \rangle dA_g,$$

where  $\phi, \psi \in H^1(M, V)$  and  $dA_g$  is the Riemannian volume on  $(\partial M, g)$ . Denote by  $I(s, t) = \langle\langle \Pi_+ u^f(t), \Pi_+ u^h(s) \rangle\rangle$ . Then, using (6), (12), and (48), we get

$$\begin{aligned} & (\partial_s^2 - \partial_t^2)I(s, t) = \langle\langle \Pi_+ u^f(t), \partial_s^2 u^h(s) \rangle\rangle - \langle\langle \Pi_+ \partial_t^2 u^f(t), u^h(s) \rangle\rangle \\ & = -\langle\langle \Pi_+ u^f(t), D(Du^h(s)) \rangle\rangle + \langle\langle D\Pi_+ u^f(t), Du^h(s) \rangle\rangle + \\ & \quad -\langle\langle D\Pi_+ u^f(t), Du^h(s) \rangle\rangle + \langle\langle D^2\Pi_+ u^f(t), u^h(s) \rangle\rangle \\ & = \int_{\partial M} (\langle \gamma(N)\Pi_+ u^f(t), D\Pi_+ u^h(s) \rangle + \langle \gamma(N)D\Pi_+ u^f(t), \Pi_+ u^h(s) \rangle) dA_g \\ & = i \int_{\partial M} \left( \langle \gamma(N)\Pi_+ \Lambda_P f(t), \Pi_- \partial_s(\Lambda_P h(s)) \rangle - \right. \\ & \quad \left. - \langle \gamma(N)\Pi_- \partial_t(\Lambda_P f)(t), \Pi_+ \Lambda_P h(s) \rangle \right) dA_g. \end{aligned} \tag{49}$$

Since  $I(0, t) = I_s(0, t) = I(s, 0) = I_t(s, 0) = 0$ , we see that  $I(s, t)$  satisfies the one-dimensional wave equation (49) in the quarter plane  $s, t > 0$  with the known right-hand side and homogeneous initial and

boundary data. Thus, we can find  $I(s, t) = \langle \Pi_+ u^f(t), \Pi_+ u^h(s) \rangle$  for any  $s, t \in \mathbb{R}_+$ .  $\square$

**Remark 4.2.** From the proof of Theorem 4.1, we obtain an explicit formula

$$(50) \quad \langle \Pi_\pm u^f(T), u^h(T) \rangle = \mathcal{L}_\pm[f, h] := \int_{\mathbb{R}} \int_{\mathbb{R}} \chi_C(t, s) J_\pm^{t,s}(f, h) ds dt.$$

Here, the quadratic form  $\mathcal{L}_\pm[\cdot, \cdot]$  is defined using the indicator function  $\chi_C$  is of the triangle  $C = \{(t, s) \in \mathbb{R}^2 : T - t \geq |T - s|\}$  and the quadratic form  $J_\pm^{t,s}(f, h)$ ,

$$(51) \quad J_\pm^{t,s}(f, h) = \frac{i}{2} \int_{\partial M} \left( \langle \gamma(N) \Pi_\pm \Lambda_P f(t), \Pi_\mp \partial_s \Lambda_P h(s) \rangle - \langle \gamma(N) \Pi_\mp \partial_t \Lambda_P f(t), \Pi_\pm \Lambda_P h(s) \rangle \right) dA_g.$$

**4.2. Global controllability.** Let  $\Sigma \subset \partial M$  be an open set and

$$X^P(\Sigma, T) = \{u^f(\cdot, T) \in L^2(M, V) : f \in PC_0^\infty(\Sigma \times (0, T), W)\},$$

where  $u^f(t)$  is the solution of (33)-(34) with the boundary condition associated with the operator  $P$ . In the following theorem,

$$\text{rad}(M, \Sigma) = \max_{x \in M} \text{dist}(x, \Sigma), \quad \text{rad}(M) = \max_{x \in M} \text{dist}(x, \partial M).$$

**Theorem 4.3.** 1. Let  $P$  be a boundary condition satisfying (18) and (19) and  $T > 2 \text{rad}(M)$ . Then the set  $X^P(\partial M, T)$  is dense in  $L^2(M, V)$ .

2. Let  $P_\Gamma$  be the local boundary condition given by (28),  $\Sigma \subset \partial M$  be an open non-empty set, and  $T > 2 \text{rad}(M, \Sigma)$ . Then the set  $X^\Gamma(\Sigma, T) = X^{P_\Gamma}(\Sigma, T)$  is dense in  $L^2(M, V)$ .

*Proof:* We start by proving claim 2. Let  $\eta \in L^2(M, V)$  be orthogonal to  $X^\Gamma(\Sigma, T)$ . Consider the dual initial boundary value problem

$$(52) \quad \begin{aligned} (i\partial_t + D)v_\eta(x, t) &= 0 \quad \text{in } M \times \mathbb{R}, \\ P_\Gamma v_\eta|_{\partial M \times \mathbb{R}} &= 0, \quad v_\eta|_{t=T} = \eta. \end{aligned}$$

Integrating by parts we see that, for any  $f \in C_0^\infty(\Sigma \times [0, T], P_\Gamma W)$ ,

$$(53) \quad \begin{aligned} 0 &= -i \langle u^f(T), v_\eta(T) \rangle \\ &= - \int_0^T \int_M (\langle i\partial_t u^f(t), v_\eta(t) \rangle - \langle u^f(t), i\partial_t v_\eta(t) \rangle) dV_g dt \\ &= \int_0^T \int_M (\langle Du^f(t), v_\eta(t) \rangle - \langle u^f(t), Dv_\eta(t) \rangle) dV_g dt \\ &= \int_0^T \int_{\partial M} \langle \gamma(N) u^f(x, t), v_\eta(x, t) \rangle dA_g(x) dt \\ &= \int_0^T \int_{\partial M} (\langle \gamma(N) P_\Gamma u^f(x, t), (I - P_\Gamma) v_\eta(x, t) \rangle + \\ &\quad + \langle \gamma(N) (I - P_\Gamma) u^f(x, t), P_\Gamma v_\eta(x, t) \rangle) dA_g dt \end{aligned}$$



$$= \int_0^T \int_{\partial M} \langle \gamma(N)f(x, t), (I - P_\Gamma)v_\eta(x, t) \rangle dA_g dt.$$

Here, the map

$$(54) \quad \mathcal{B} : L^2(M, V) \mapsto \mathcal{D}'(\partial M \times \mathbb{R}, W), \quad \mathcal{B}(\eta) = v_\eta|_{\partial M \times \mathbb{R}},$$

is continuous. This may be shown by an approximation of  $\eta \in L^2(M, V)$  by  $C_0^\infty(M, V)$  sections and applying the uniform boundedness principle for the distributions. Since equation (53) is valid for any  $f \in C_0^\infty(\Sigma \times [0, T], P_\Gamma W)$ , we see that  $(I - P_\Gamma)v_\eta|_{\Sigma \times (0, T)} = 0$ . Together with the boundary condition of (52), this makes it possible to apply Theorem 3.1. Hence  $v_\eta$  vanishes in  $K_{\Sigma, T/2}$ . However, the set  $K_{\Sigma, T/2}$  contains an open neighborhood of  $M \times \{\frac{T}{2}\}$ . As also  $P_\Gamma v_\eta|_{\partial M \times \mathbb{R}} = 0$ , this implies that  $v_\eta = 0$  in  $M \times \mathbb{R}$ . In particular,  $\eta = v_\eta(T) = 0$  proving claim 2.

Claim 1. may be proven by the same arguments as claim 2. by taking  $P$  instead of  $P_\Gamma$  and using property (18).  $\square$

As can be seen in the above proof, for any  $t_0$  and  $f \in C_0^\infty(\partial M \times (0, t_0), P_\Gamma W)$ , and  $v_\eta$  solving (52) with  $\eta \in L^2(M, V)$ ,

$$(55) \quad \langle \langle u^f(t_0), v_\eta(t_0) \rangle \rangle = i \int_0^{t_0} \int_{\partial M} \langle \gamma(N)f(x, t), v_\eta(x, t) \rangle dA_g dt,$$

where the right hand side is understood as a distribution pairing.

**4.3. Local controllability results.** When  $P = P_\Gamma$  is the local projection on  $\partial M$ , we define

$$\begin{aligned} X_+^\Gamma(\Sigma, T) &= \{\Pi_+ u^f(\cdot, T) \in L^2(M, V_+) : f \in C_0^\infty(\Sigma \times (0, T), P_\Gamma W)\}, \\ X_-^\Gamma(\Sigma, T) &= \{\Pi_- u^f(\cdot, T) \in L^2(M, V_-) : f \in C_0^\infty(\Sigma \times (0, T), P_\Gamma W)\}. \end{aligned}$$

For an arbitrary set  $S \subset M$ , we also denote

$$L^2(S, V) = \{v \in L^2(M, V) : \text{supp}(v) \subset \overline{S}\}.$$

**Theorem 4.4.** *Let  $P_\Gamma$  be the local projection on  $\partial M$  given by (28). Then, for any  $T > 0$  and open  $\Sigma \subset \partial M$ , the sets  $X_\pm^\Gamma(\Sigma, T)$  are dense in the subspaces  $\Pi_\pm L^2(M(\Sigma, T), V) \subset L^2(M, V)$ , correspondingly.*

*Proof:* We will prove this result for  $X_+^\Gamma(\Sigma, T)$ , the claim for  $X_-^\Gamma(\Sigma, T)$  may be obtained in the same manner.

Let  $\eta \in \Pi_+ L^2(M(\Sigma, T), V)$  satisfy

$$\langle \langle \eta, \Pi_+ u^f(T) \rangle \rangle = 0 \quad \text{for any } f \in C_0^\infty(\Sigma \times (0, T), P_\Gamma W).$$

Consider the initial boundary value problem

$$(56) \quad \begin{aligned} (i\partial_t + D)v_\eta(x, t) &= 0 \quad \text{in } M \times \mathbb{R}, \\ P_\Gamma v_\eta|_{\partial M \times \mathbb{R}} &= 0, \\ \Pi_+ v_\eta|_{t=T} &= \eta, \quad \Pi_- v_\eta|_{t=T} = 0. \end{aligned}$$

Then  $v_\eta \in C(\mathbb{R}, L^2(M, V))$  and, as noted in the proof of Theorem 4.3, it has a well defined trace  $v_\eta|_{\partial M \times \mathbb{R}} \in \mathcal{D}'(\partial M \times \mathbb{R})$ . Same considerations as in (53) show that

$$0 = \int_0^T \int_{\partial M} \langle \gamma(N)f(x, t), (I - P_\Gamma)v_\eta(x, t) \rangle dA_g dt,$$

for any  $f \in C_0^\infty(\Sigma \times [0, T], P_\Gamma W)$ . Therefore,  $(I - P_\Gamma)v_\eta|_{\Sigma \times (0, T)} = 0$ . Together with the boundary condition in (56), this imply that  $v_\eta|_{\Sigma \times (0, T)} = 0$ , i.e.  $\Pi_\pm v_\eta|_{\Sigma \times (0, T)} = 0$ . Using (46) and the fact that  $\Pi_- v_\eta|_{t=T} = 0$ , we see that  $\Pi_- v_\eta(T+s) = -\Pi_- v_\eta(T-s)$  and  $\Pi_+ v_\eta(T+s) = \Pi_+ v_\eta(T-s)$ . Therefore,

$$(57) \quad \Pi_\pm v_\eta|_{\Sigma \times (T, 2T)} = 0 \quad \text{and} \quad \text{supp}(v_\eta|_{\Sigma \times (0, 2T)}) \subset \Sigma \times \{T\}.$$

We intend to show that  $v_\eta|_{\partial M \times (0, 2T)} = 0$ . Due to the boundary condition in (56) this is equivalent to

$$(58) \quad \int_0^{2T} \int_\Sigma \langle \gamma(N)f(x, t), v_\eta(x, t) \rangle dA_g dt = 0$$

for any  $f \in C_0^\infty(\Sigma \times (0, 2T), P_\Gamma W)$ . In view of (57), it is enough to prove that

$$(59) \quad \begin{aligned} & \int_{T-2\varepsilon}^{T+2\varepsilon} \int_\Sigma \langle \gamma(N)f_\varepsilon(x, t), v_\eta(x, t) \rangle dA_g dt \\ &= i \langle \langle u^{f_\varepsilon}(T+2\varepsilon), v_\eta(T+2\varepsilon) \rangle \rangle \longrightarrow 0, \quad \text{when } \varepsilon \rightarrow 0. \end{aligned}$$

Here  $f_\varepsilon(\cdot, t) = \chi_\varepsilon(t-T)f(\cdot, t)$  and  $\chi_\varepsilon(s)$  is a smooth cut-off function,  $\chi(s) = 1$  for  $s \in (-1, 1)$ ,  $\text{supp}(\chi) \subset (-2, 2)$  with  $\chi_\varepsilon(s) = \chi(s/\varepsilon)$ . Note that the equality in (59) follows from (55).

Assume, in the beginning, that  $f_\varepsilon(x, t) = \chi_\varepsilon(t-T)F_\partial(x)$ , where  $F_\partial = F|_{\partial M}$ ,  $F \in C^\infty(M, V)$ . We can represent the solution as

$$u^{f_\varepsilon}(x, t) = \chi_\varepsilon(t-T)F(x) + \omega_\varepsilon(x, t),$$

where  $\omega_\varepsilon$  is the solution to

$$\begin{aligned} (i\partial_t + D)\omega_\varepsilon &= -(i\partial_t + D)(\chi_\varepsilon F) = -\frac{i}{\varepsilon}\chi'((t-T)/\varepsilon)F - \chi_\varepsilon(t-T)DF, \\ P_\Gamma \omega_\varepsilon|_{\partial M \times \mathbb{R}} &= 0, \quad \omega_\varepsilon|_{t=T-2\varepsilon} = 0. \end{aligned}$$

We represent  $F$  and  $DF$  as  $L^2(M)$ -converging series of eigenfunctions  $\phi_k$  of  $D_\Gamma$ ,

$$F = \sum_{k=1}^{\infty} a_k \phi_k, \quad DF = \sum_{k=1}^{\infty} b_k \phi_k.$$

Then

$$(60) \quad \omega_\varepsilon(T+t) = \sum_{k=1}^{\infty} (w_k^1(t) + w_k^2(t)) \phi_k = \omega_\varepsilon^1(T+t) + \omega_\varepsilon^2(T+t),$$

$$w_k^1(t) = \frac{a_k}{i\varepsilon} \int_{T-2\varepsilon}^t e^{i\lambda_k(t-s)} \chi'(s/\varepsilon) ds, \quad w_k^2(t) = -b_k \int_{T-2\varepsilon}^t e^{i\lambda_k(t-s)} \chi(s/\varepsilon) ds.$$

It then follows from (60) that

$$(61) \quad \|\omega_\varepsilon^1(T+2\varepsilon)\|_{L^2} \leq C_1 \|F\|_{L^2}, \quad \|\omega_\varepsilon^2(T+2\varepsilon)\|_{L^2} \leq C_2 \varepsilon \|DF\|_{L^2}.$$

What is more, introducing

$$\omega_\varepsilon^{1,N}(T+t) = \sum_{|\lambda_k| > N} w_k^1(t) \phi_k,$$

we see that

$$(62) \quad \lim_{N \rightarrow \infty} \|\omega_\varepsilon^{1,N}(T+2\varepsilon)\|_{L^2} = 0,$$

uniformly with respect to  $\varepsilon \in (0, \varepsilon_0)$ . To estimate  $\omega_\varepsilon^1 - \omega_\varepsilon^{1,N}$ , use integration by parts in the equation for  $w_k^1(t)$  in (60) to get

$$(63) \quad |w_k^1(2\varepsilon)| \leq C_3 N |a_k| \varepsilon \quad \text{when } |\lambda_k| \leq N.$$

Combining (62)–(63) we see that  $\|\omega_\varepsilon^1(T+2\varepsilon)\|_{L^2} \rightarrow 0$  when  $\varepsilon \rightarrow 0$ . This and the estimate for  $\omega_\varepsilon^2(T+2\varepsilon)$  in (61) imply the desired result (59) for  $f_\varepsilon(x, t) = \chi_\varepsilon(t-T)F_\partial(x)$ .

Returning to a general  $f \in C_0^\infty(\Sigma \times (0, 2T), P_\Gamma W)$ , it remains to estimate  $\omega_\varepsilon(T+2\varepsilon)$  for  $f_\varepsilon(x, t) = (t-T)\chi_\varepsilon(t-T)\tilde{f}(x, t)$  with smooth  $\tilde{f}(x, t)$ . Arguments similar to those leading to the estimate of  $\omega_\varepsilon^2(T+2\varepsilon)$  in (61) show that  $\|\omega_\varepsilon(T+2\varepsilon)\|_{L^2} \rightarrow 0$  when  $\varepsilon \rightarrow 0$ . This proves (58) and, therefore, Theorem 4.4.  $\square$

We complete this section by describing a procedure to find  $\text{rad}(M, \Sigma)$ .

**Corollary 4.5.** *For any open  $\Sigma \subset \partial M$ , the Cauchy data set  $C_0(D)$  and the induced bundle structure on  $\partial M$  determine  $\text{rad}(M, \Sigma)$ .*

*Proof:* By the local controllability result, Theorem 4.4 we see that, if  $T_1 > \text{rad}(M, \Sigma)$ , then the following claim (A) is valid:

(A) For any  $T_2 > 0$  and  $f \in C_0^\infty(\partial M \times \mathbb{R}_+, P_\Gamma W)$  there are  $h_j^+, h_j^- \in C_0^\infty(\Sigma \times (0, T_1), P_\Gamma W)$ ,  $j = 1, 2, \dots$  such that

$$(64) \quad \begin{aligned} \lim_{j \rightarrow \infty} \|\Pi_+(u^{h_j^+}(T_1) - u^f(T_2))\|_{L^2} &= 0, \\ \lim_{j \rightarrow \infty} \|\Pi_-(u^{h_j^-}(T_1) - u^f(T_2))\|_{L^2} &= 0. \end{aligned}$$

On the other hand, if  $T_1 < \text{rad}(M, \Sigma)$ , we see that for  $T_2 > \text{rad}(M, \Sigma)$  there is  $f$  such that  $\text{supp}(u^f(T_2)) \not\subset M(\Sigma, T_1)$ . Therefore, claim (A) is not valid.

As by Theorem 4.1 the norms on the left-hand sides of (64) can be evaluated using the boundary data, we find  $\text{rad}(M, \Sigma)$  by taking the infimum of all  $T_1 > 0$  such that (A) is valid.  $\square$

## 5. GENERALIZED BOUNDARY SOURCES

**5.1. The wave norm.** We define the wave operators

$$\mathcal{U}^T : PC_0^\infty(\partial M \times (0, T), W) \rightarrow L^2(M, V), \quad \mathcal{U}^T f = u^f(T),$$

where  $T = 3 \text{ rad}(M)$  and  $u^f(t) = u^f(x, t)$  is the solution of (33)-(34). By means of the wave operators, we define a semi-norm, or the wave semi-norm on the space of boundary sources as

$$(65) \quad \|f\|_{\mathcal{F}} = \|\mathcal{U}^T f\|_{L^2(M, V)}.$$

By Theorem 4.1, the knowledge of the operator  $\Lambda_P$  and the induced bundle structure (31) on  $\partial M$  enables us to calculate explicitly the semi-norm (65) of any  $f \in PC_0^\infty(\partial M \times (0, T), W)$ .

To complete the space of boundary sources, we define the equivalence relation  $\sim$  on this space by setting

$$f \sim h \text{ if and only if } u^f(T) = u^h(T).$$

Further, we define the space  $\mathcal{B}$  as

$$\mathcal{B} = PC_0^\infty(\partial M \times (0, T), W) / \sim.$$

Then (65) becomes a norm on  $\mathcal{B}$ . Finally, we complete  $\mathcal{B}$  with respect to the norm (65). This space, denoted by  $\mathcal{F} = \overline{\mathcal{B}}$ , consists of the sequences of sources which are the Cauchy sequences with respect to the norm (65). We denote these sequences by  $\hat{f}$ ,

$$\hat{f} = (f_j)_{j=0}^\infty, \quad f_j \in PC_0^\infty(\partial M \times (0, T), W).$$

Also, we denote  $J : PC_0^\infty(\partial M \times (0, T), W) \rightarrow \mathcal{F}$  the factor-projection that maps  $f$  to its equivalence class in  $\mathcal{B}$ .

The sources  $\hat{f} \in \mathcal{F}$  are called *generalized sources* in the sequel. The corresponding waves are denoted by

$$(66) \quad \mathcal{W}\hat{f}(t) = \lim_{j \rightarrow \infty} u^{f_j}(t) \quad \text{for } t \geq T,$$

where the convergence in the right-hand side is the  $L^2(M, V)$ -convergence. By the global controllability, Theorem 4.3 and the definition (65) of the norm, the map  $\hat{f} \mapsto \mathcal{W}\hat{f}(t)$  is an isometry,  $\mathcal{F} \rightarrow L^2(M, V)$ . We extend the notation  $\mathcal{W}\hat{f}(t)$  for all  $t \in \mathbb{R}$  by defining

$$(67) \quad \mathcal{W}\hat{f}(t) = \exp(i(t - T)D_P)\mathcal{W}\hat{f}(T) \quad \text{for } t \in \mathbb{R}.$$

Due to this formula we call  $\mathcal{W}\hat{f}(t)$  a *generalized wave*.

**Remark 5.1.** The choice of the parameter  $T = 3 \text{ rad}(M)$  in the above construction can be replaced with any fixed  $T > 2 \text{ rad}(M)$ .

We note that the above construction of the space of generalized sources is well-known in PDE-control, e.g. [40, 50].

For  $t \in \mathbb{R}$  and any boundary source  $\hat{f} \in \mathcal{F}$ , we can find if the condition

$$(68) \quad \partial_t^l \mathcal{W}\hat{f}(t) \in L^2(M, V)$$

is satisfied for a given  $l \in \mathbb{Z}_+$ . Indeed, for  $l = 1$ , condition (68) for  $\hat{f} = (f_j)_{j=1}^\infty \in \mathcal{F}$  is equivalent to the existence of  $\hat{h} \in \mathcal{F}$  such that

$$(69) \quad \begin{aligned} & \lim_{\delta \rightarrow 0} \left\| \frac{\mathcal{W}f(t) - \mathcal{W}f(t - \delta)}{\delta} - \mathcal{W}h(t) \right\|_{L^2(M, V)} \\ &= \lim_{\delta \rightarrow 0} \left\| \mathcal{W} \left( \frac{\hat{f} - \mathcal{T}_\delta \hat{f}}{\delta} - \hat{h} \right) (t) \right\|_{L^2(M, V)} = 0, \end{aligned}$$

$t > T$ . Here  $\mathcal{T}_\delta : f(x, t) \mapsto f(x, t - \delta)$  is the time-shift naturally extended from  $PC_0^\infty(\partial M \times (0, T), W)$  to  $\mathcal{F}$ . Note that the validity of (69) is independent of value of  $t > T$  as the norms  $\|\mathcal{W}\hat{f}(t)\|_{L^2}$  are independent of  $t$ . Using the induced bundle structure on  $\partial M$  and the response operator  $\Lambda_P$  we can verify if a given generalized source  $\hat{f} = (f_j)_{j=1}^\infty$  and a trial  $\hat{h} = (h_j)_{j=1}^\infty$  satisfy condition (69). In this case we denote

$$\hat{h} = \partial_t \hat{f} \quad \text{and} \quad \hat{f} \in \mathcal{D}(\partial_t).$$

Having defined  $\partial_t^{k-1} \hat{f}$  we define  $\partial_t^k \hat{f}$ , if it exists, by induction. The spaces of generalized sources which have  $k \in \mathbb{Z}_+$  time derivatives will be denoted by  $\mathcal{F}^k = \mathcal{D}(\partial_t^k)$  with  $\mathcal{F}^\infty = \bigcap_{k \in \mathbb{Z}_+} \mathcal{F}^k$ . Clearly,

$$\partial_t Jf = J(f_t), \quad f \in PC_0^\infty(\partial M \times (0, T), W),$$

that is,  $\partial_t$  is an extension of the time derivative  $f \mapsto f_t$  defined in the usual way for  $f \in PC_0^\infty(\partial M \times (0, T), W)$ . Note that by (67) and (68),

$$(70) \quad \hat{f} \in \mathcal{D}(\partial_t^k), \quad \text{if and only if} \quad \mathcal{W}\hat{f}(t) \in \mathcal{D}(D_P^k), \quad t \in \mathbb{R}.$$

In addition,

$$(71) \quad \mathcal{W}(\partial_t^k \hat{f})(t) = i^k D_P^k \mathcal{W}\hat{f}(t) \quad \text{for } \hat{f} \in \mathcal{F}^k.$$

**5.2. Boundary sources corresponding to eigenfunctions.** When  $\hat{f} \in \mathcal{F}^1$  we can evaluate, using Theorem 4.1, the inner products

$$\begin{aligned} \langle \langle D_P \mathcal{W}\hat{f}(T), \mathcal{W}\hat{f}(T) \rangle \rangle &= -i \langle \langle \mathcal{W}(\partial_t \hat{f})(T), \mathcal{W}\hat{f}(T) \rangle \rangle, \\ \langle \langle D_P \mathcal{W}\hat{f}(T), D_P \mathcal{W}\hat{f}(T) \rangle \rangle &= \langle \langle \mathcal{W}(\partial_t \hat{f})(T), \mathcal{W}(\partial_t \hat{f})(T) \rangle \rangle, \end{aligned}$$

as well as the inner products  $\langle \langle \mathcal{W}\hat{f}(T), \mathcal{W}\hat{f}(T) \rangle \rangle$ . On the other hand, by the Hilbert-Courant min-max principle the eigenvalues  $[\lambda_j]^2$  of  $D_P^2$  can be found as

$$(72) \quad [\lambda_j]^2 = \sup_{u_1, \dots, u_{n-1}} \inf_{u_n \perp u_1, \dots, u_{n-1}} \frac{\langle \langle D_P u_j, D_P u_j \rangle \rangle}{\langle \langle u_j, u_j \rangle \rangle},$$

where  $u_j$  are in  $\mathcal{D}(D_P)$ . This makes it possible to prove the following result:

**Proposition 5.2.** *The induced bundle structure (31) on  $\partial M$  and the response operator  $\Lambda_P$  determine the eigenvalues  $\lambda_j$  of  $D_P$  and the generalized sources  $\hat{h}_j \in \mathcal{F}^\infty$ ,  $j = 1, 2, \dots$ , such that  $\phi_j = \mathcal{W}\hat{h}_j(T)$  are orthonormal eigenfunctions of  $D_P$  that form a complete basis of  $L^2(M, V)$ .*

*Proof:* Using (72), we can find all the eigenvalues of  $D_P^2$ . Indeed, by the global controllability, Theorem 4.3 and relation (71),  $\mathcal{W}\hat{f}(T)$  runs over  $\mathcal{D}(D_P)$  when  $\hat{f}$  runs over  $\mathcal{F}^1$ . At the same time, the minimizers in (72) are those generalized sources  $\hat{f}_j \in \mathcal{F}^1$  that the waves  $u_j = \mathcal{W}\hat{f}_j(T)$  are the corresponding orthonormal eigenvectors of  $D_P^2$ . Furthermore, since we can evaluate  $\langle\langle D_P u_j, u_j \rangle\rangle$ , we can decompose the eigenspace of  $D_P^2$ , which corresponds to an eigenvalue  $[\lambda_j]^2$ , into a direct sum of the eigenspaces of  $D_P$  which corresponds to the eigenvalues  $\lambda_j$  and  $-\lambda_j$ . Hence, we can find  $\hat{h}_j \in \mathcal{F}^1$  such that  $\phi_j = \mathcal{W}\hat{h}_j(T)$  are the orthonormal eigenfunctions of  $D_P$ . Clearly,  $\hat{h}_j \in \mathcal{F}^\infty$ .  $\square$

Applying Theorem 4.1 to  $\mathcal{W}\hat{f}(t)$  and  $\mathcal{W}\hat{h}_j(T)$ , we can evaluate the Fourier coefficients  $w_j(t) = \langle\langle \mathcal{W}\hat{f}(t), \phi_j \rangle\rangle$ ,  $j = 1, 2, \dots$ , of  $\mathcal{W}\hat{f}(t)$ . Using  $w_j(t)$ , we can compute the norms  $\|(1 + |D_P|)^s \mathcal{W}\hat{f}(t)\|_{L^2(M)}$ ,  $s \geq 0$ . We define the spaces  $\mathcal{F}^s$  as the spaces of those generalized sources  $\hat{f} \in \mathcal{F}$  for which  $\mathcal{W}\hat{f}(t) \in \mathcal{D}(|D_P|^s)$  and define

$$\|\hat{f}\|_{\mathcal{F}^s} = \|(1 + |D_P|)^s \mathcal{W}\hat{f}(t)\|_{L^2(M)} = \sum_{j=1}^{\infty} (1 + |\lambda_j|)^{2s} |w_j(t)|^2,$$

where the right-hand side is independent of  $t \in \mathbb{R}$ .

## 6. INDEX FORMULAE USING BOUNDARY DATA

In this section we use boundary data to verify if the decomposition condition (24) is valid and give formulae for the Fredholm index of the operator  $D_P^+$  appearing in decomposition (25).

**6.1. Decomposition of  $D_P$ .** Let us consider first the Dirac operator  $D_\Gamma$  associated with  $P_\Gamma$ . Using properties (8) of the chirality operator, we can see that, if  $u \in \mathcal{D}(D_\Gamma)$  and  $\Pi_+ u \in \mathcal{D}(D_\Gamma)$ , then  $u|_{\partial M} = 0$ . Therefore, the decomposition condition (24) is not satisfied for the Dirac-type operator  $D_\Gamma$ . However, there is a large number of important examples of projectors  $P$  when the decomposition condition (24) is valid for the corresponding Dirac-type operator  $D_P$ . Some of these are discussed below.

**6.2. Decomposition of  $\mathcal{F}$ .** Assume that we are given the operator  $\Lambda_P$  with a possibly non-local boundary operator  $P$  and the induced bundle structure on  $\partial M$ . Then the formula (50) determines the non-negative sesquilinear forms  $\mathcal{L}_+[\cdot, \cdot]$  and  $\mathcal{L}_-[\cdot, \cdot]$  on  $PC_0^\infty(\partial M \times (0, T), W)$  such

that

$$\mathcal{L}_\pm(f, h) = \langle \Pi_\pm u^f(T), u^h(T) \rangle.$$

Observe that the quadratic form  $\mathcal{L} = \mathcal{L}_+ + \mathcal{L}_-$  defines a semi-norm  $\sqrt{\mathcal{L}[f, f]}$  that coincides, on  $PC_0^\infty(\partial M \times (0, T), W)$ , with the semi-norm defined by (65)

$$(73) \quad \mathcal{L}[f, f] = \mathcal{L}_+[f, f] + \mathcal{L}_-[f, f] = \|f\|_{\mathcal{F}}^2.$$

Therefore,  $\mathcal{L}_+$  and  $\mathcal{L}_-$  can be extended to  $\mathcal{F}$ .

In turn, these forms give rise to two bounded non-negative operators  $B_\pm$  on  $\mathcal{F}$  so that

$$\mathcal{L}_\pm[\hat{f}, \hat{h}] = (B_\pm \hat{f}, \hat{h})_{\mathcal{F}}.$$

The fact that  $\Pi_+ L^2(M, V) \oplus \Pi_- L^2(M, V) = L^2(M, V)$  implies that  $B_\pm$  are orthoprojections in  $\mathcal{F}$  satisfying  $B_+ \oplus B_- = I$ . Introducing an operator  $S$ ,

$$S\hat{f} = \mathcal{W}\hat{f}(0), \quad \hat{f} \in \mathcal{F},$$

we have

$$(74) \quad SB_\pm \hat{f} = \Pi_\pm S\hat{f}.$$

**6.3. Index formulae.** At first we prove Theorem 2.6.

*Proof:* Using the operator  $S$ , equation (71) may be written as

$$(75) \quad D_P = S(-i\partial_t)S^{-1}.$$

Together with (74), this equation imply that the condition  $\Pi_\pm \mathcal{D}(D_P) \subset \mathcal{D}(D_P)$  is satisfied if and only if  $B_\pm \mathcal{D}(\partial_t) \subset \mathcal{D}(\partial_t)$ . This proves claim 1.

If decomposition (24) is valid, so that  $D_P$  can be decomposed as in (25), formula (75) implies that  $\partial_t$  can be decomposed as

$$\partial_t = \begin{pmatrix} 0 & \partial_t^- \\ \partial_t^+ & 0 \end{pmatrix} : B_+ \mathcal{D}(\partial_t) \oplus B_- \mathcal{D}(\partial_t) \rightarrow B_- \mathcal{F} \oplus B_+ \mathcal{F},$$

where  $\partial_t^\pm$  are the restrictions of  $\partial_t$  on  $B_\pm \mathcal{D}(\partial_t)$ , correspondingly. As  $D_P^\pm$  are Fredholm operators, we conclude from (74) and (75) that  $\partial_t^\pm$  are Fredholm operators and  $\text{Ind}(\partial_t^+) = \text{Ind}(D_P^+)$ . This proves claim 2.  $\square$

**Corollary 6.1.** *Assume that condition (24) is valid for  $D_P$ . Then*

$$\text{Ind}(D_P^+) = \text{sgn}([B_+ - B_-]|_{\text{Ker}(\partial_t)}).$$

*Proof:* As  $FD + DF = 0$ , we have  $F : \text{Ker}(D_P) \rightarrow \text{Ker}(D_P)$  so that  $\Pi_\pm : \text{Ker}(D_P) \rightarrow \text{Ker}(D_P)$ . Thus,  $\text{Ind}(\partial_t^+)$  is equal to

$$\dim \text{Ker}(\partial_t|_{B_+ \mathcal{F}^1}) - \dim \text{Ker}(\partial_t|_{B_- \mathcal{F}^1}) = \text{sgn}([B_+ - B_-]|_{\text{Ker}(\partial_t)}),$$

where we use the fact that  $B_\pm$  are orthoprojectors in  $\mathcal{F}$ .  $\square$

By Proposition 5.2, using the boundary data it is possible to find generalized sources  $\hat{h}_j = (h_j^k)_{k=1}^\infty \in \mathcal{F}^\infty$  such that  $\mathcal{W}\hat{h}_j(T) = \phi_j$ , with  $\{\phi_j\}_{j=1}^\infty$  being an orthonormal basis of the eigenfunctions of  $D_P$ . Let  $\phi_j$ ,  $j = 1, \dots, \nu$ , where  $\nu = \dim \text{Ker}(D_P)$ , be the eigenfunctions spanning  $\text{Ker}(D_P)$ . Then, for any  $t_0 > T$ ,

$$\lim_{k \rightarrow \infty} \Lambda_P(h_j^k)|_{\partial M \times (t_0, \infty)} = 1(t) \phi_j(x)|_{\partial M}, \quad j = 1, \dots, \nu,$$

with  $1(t) \equiv 1$ . Thus,  $\Lambda_P$  determines the subspace

$$\mathcal{N} = \text{span}\{\phi_j|_{\partial M} : j = 1, \dots, \nu\} \subset (1 - P)C^\infty(\partial M, W).$$

As  $F : \mathcal{N} \rightarrow \mathcal{N}$  and  $F = \Pi_+ - \Pi_-$ , Corollary 6.1 implies the following result:

**Lemma 6.2.** *Assume that  $D_P$  satisfies condition (24). Then*

$$\text{Ind}(D_P^+) = \text{sgn}(F|_{\mathcal{N}}),$$

where the right-hand side can be evaluated in terms of the induced bundle structure (31) on  $\partial M$  and the response operator  $\Lambda_P$ .

**Example 1, continued.** It follows from the definitions of  $\mathcal{F}$  and the relative and absolute boundary conditions, that the Dirac operators  $(d + \delta)_r$  and  $(d + \delta)_a$  can be decomposed into  $(d + \delta)_r^\pm$  and  $(d + \delta)_a^\pm$ . Clearly,  $\text{Ker}[(d + \delta)_r^+]$  consists of the relative harmonic forms of the even order and  $\text{Ker}[(d + \delta)_r^-]$  of relative harmonic forms of the odd order. Similarly,  $\text{Ker}[(d + \delta)_a^+]$  and  $\text{Ker}[(d + \delta)_a^-]$  consists of absolute harmonic forms of an even and odd order, correspondingly. Therefore,  $\text{Ind}[(d + \delta)_r^+]$  is the Euler characteristic  $\chi(M)$  of  $M$ . By the Poincaré duality,  $\text{Ind}[(d + \delta)_a^+] = \pm \text{Ind}[(d + \delta)_r^+]$  depending on  $n = \dim(M)$ . Thus, Theorem 2.6 gives a representation of the Euler characteristic  $\chi(M)$  in terms of the Cauchy data set  $C_0(D)$  of the hyperbolic form-Dirac equation.

**Example 2, continued.** Similarly to example 1, the index of the extended Maxwell operator  $i(d - \delta_a)$  with the electric boundary condition is the Euler characteristic  $\chi(M)$ .

**Example 3, continued.** The generalized signature operator on even dimensional manifold with boundary satisfies decomposition condition (24). This was first considered in the famous paper [1] (for further developments in this area see e.g. [12, 19, 18, 41, 43]). Theorem 2.6 provides a possibility to find the index of the signature operator from the Cauchy data of its Dirac operator.

## 7. RECONSTRUCTION OF THE MANIFOLD

In this section we will show how to reconstruct the manifold,  $M$  and metric,  $g$  from the response operator  $\Lambda_\Gamma$ . To this end, we will show that the boundary data determine the set of the *boundary distance functions* that determine  $(M, g)$  up to an isometry.



**7.1 Controlling supports of generalized waves.** We start by fixing certain notations. Let

$$(76) \quad T_1 = T + 4 \sup \{ \text{rad}(M, \Sigma) : \Sigma \neq \emptyset \}.$$

Then  $T_1$  is determined by the boundary data and  $T_1 - T > \text{diam}(M)$ .

Let  $\Sigma_j \subset \partial M$  be non-empty open disjoint sets,  $1 \leq j \leq J$  and  $\tau_j^-$  and  $\tau_j^+$  be positive times with

$$0 < \tau_j^- < \tau_j^+ \leq T_1 - T, \quad 1 \leq j \leq J.$$

Let  $S = S(\{\Sigma_j, \tau_j^-, \tau_j^+\}_{j=1}^J) \subset M$  be the intersection of slices,

$$(77) \quad S = \bigcap_{j=1}^J (M(\Sigma_j, \tau_j^+) \setminus M(\Sigma_j, \tau_j^-)).$$

Our first goal is to find, using the boundary data, whether the set  $S$  contains an open ball or not. To this end, we give the following definition.

**Definition 7.1.** The set  $Z = Z(\{\Sigma_j, \tau_j^-, \tau_j^+\}_{j=1}^J)$  is the set of the generalized sources  $\hat{f} \in \mathcal{F}^\infty$  that generate the waves  $\mathcal{W}\hat{f}(t)$  with

- (1)  $\text{supp}(\mathcal{W}\hat{f}(T_1)) \subset M(\Sigma_j, \tau_j^+)$  for all  $j$ ,  $1 \leq j \leq J$ ,
- (2)  $\mathcal{W}\hat{f}(T_1) = 0$  in  $M(\Sigma_j, \tau_j^-)$  for all  $j$ ,  $1 \leq j \leq J$ .

**Theorem 7.2.** *Given the induced bundle structure on  $\partial M$  and the response operator  $\Lambda_\Gamma$ , we can determine whether a given boundary source  $\hat{f} \in \mathcal{F}^\infty$  is in  $Z$  or not.*

*Proof:* Let  $\hat{f} = (f_k)_{k=0}^\infty \in \mathcal{F}^\infty$  be a generalized source. Consider first the question whether  $\text{supp}(\mathcal{W}\hat{f}(T_1)) \subset M(\Sigma_j, \tau_j^+)$  or equivalently,  $\text{supp}(\Pi_+ \mathcal{W}\hat{f}(T_1)) \subset M(\Sigma_j, \tau_j^+)$  and  $\text{supp}(\Pi_- \mathcal{W}\hat{f}(T_1)) \subset M(\Sigma_j, \tau_j^+)$ . By Theorem 4.4 this is equivalent to the existence of generalized sources,

$$\hat{h}^+ = (h_\ell^+)_{\ell=0}^\infty \text{ and } \hat{h}^- = (h_\ell^-)_{\ell=0}^\infty, \quad h_\ell^\pm \in C_0^\infty(\Sigma_j \times (0, \tau_j^+), P_\Gamma W),$$

such that

$$(78) \quad \lim_{k, \ell \rightarrow \infty} \|\Pi_+(u^{f_k}(T_1) - u_t^{h_\ell^+}(\tau_j^+))\|_{L^2(M, V)} = 0,$$

$$(79) \quad \lim_{k, \ell \rightarrow \infty} \|\Pi_-(u^{f_k}(T_1) - u_t^{h_\ell^-}(\tau_j^+))\|_{L^2(M, V)} = 0.$$

By Theorem 4.1, we can evaluate the left-hand sides of (78) and (79) using the induced bundle structure (31) on  $\partial M$  and the operator  $\Lambda_\Gamma$ . Thus, for a given  $\hat{f}$ , we can verify if condition 1. is satisfied.

Next, consider condition 2. As

$$(i\partial_t + D)\mathcal{W}\hat{f} = 0 \text{ in } M \times \mathbb{R}, \quad P_\Gamma(\mathcal{W}\hat{f}|_{\partial M \times \mathbb{R}}) = 0,$$

condition 2. implies, due to the finite propagation speed, that

$$\mathcal{W}\hat{f} = 0 \text{ in } K_j = \{(x, t) \in M \times \mathbb{R} : \text{dist}(x, \Sigma_j) + |t - T_1| < \tau_j^-\},$$

for all  $j = 1, \dots, J$ . In particular,

$$(80) \quad \mathcal{W}\hat{f}|_{\Sigma_j \times (T_1 - \tau_j^-, T_1 + \tau_j^-)} = \lim_{j \rightarrow \infty} \Lambda_\Gamma f_j|_{\Sigma_j \times (T_1 - \tau_j^-, T_1 + \tau_j^-)} = 0.$$

On the other hand, if (80) takes place, then Theorem 3.1 implies that  $\mathcal{W}\hat{f} = 0$  in  $K_j$ . As condition (80) can be verified given the induced bundle structure on  $\partial M$  and the response operator  $\Lambda_\Gamma$ , we can verify condition 2.  $\square$

**Theorem 7.3.** *Let  $S$  and  $Z$  be as above. The following alternative holds:*

1. *If  $S$  contains an open ball, then  $\dim(Z) = \infty$ ,*
2. *If  $S$  does not contain an open ball, then  $Z = \{0\}$ .*

*Proof:* 1. Let  $B \subset S$  be an open ball and  $0 \neq \varphi \in C_0^\infty(B, V)$ . By the global controllability, Theorem 4.3, there is  $\hat{f} \in \mathcal{F}^\infty$  such that

$$\mathcal{W}\hat{f}(T_1) = \varphi.$$

As  $C_0^\infty(B, V) \subset \mathcal{D}(D_\Gamma^\infty)$ , then  $\hat{f} \in Z$ . As  $\phi \in C_0^\infty(B, V) \setminus \{0\}$  is arbitrary, this implies that  $\dim(Z) = \infty$ .

2. Assume  $S$  does not contain an open ball and suppose that there is  $0 \neq \hat{f} \in Z$ . Then, by conditions 1., 2. in Definition 7.1,

$$\text{supp}(\mathcal{W}\hat{f}(T_1)) \subset \bigcap_{j=1}^J M(\Sigma_j, \tau_j^+) = S^+,$$

$$\mathcal{W}\hat{f}(x, T_1) = 0, \quad \text{for } x \in \bigcup_{j=1}^J M(\Sigma_j, \tau_j^-) = S^-.$$

Thus  $\text{supp}(\mathcal{W}\hat{f}(T_1)) \subset S^+ \setminus S^-$ . As  $S$  does not contain a ball, then  $S^+ \setminus S^-$  is nowhere dense. Since  $\mathcal{W}\hat{f}(T_1)$  is smooth, it vanishes everywhere in  $M$ . This is a contradiction with the assumption that  $\hat{f} \neq 0$  in  $\mathcal{F}$ .  $\square$

We are now ready to construct the set of the boundary distance functions. For each  $x \in M$ , the corresponding boundary distance function,  $r_x \in C(\partial M)$  is given by

$$r_x : \partial M \rightarrow \mathbb{R}_+, \quad r_x(z) = \text{dist}(x, z), \quad z \in \partial M.$$

The boundary distance functions define *the boundary distance map*  $\mathcal{R} : M \rightarrow C(\partial M)$ ,  $\mathcal{R}(x) = r_x$ , which is continuous and injective (see [34, 29]). Denote by

$$\mathcal{R}(M) = \{r_x \in C(\partial M) : x \in M\}$$

the image of  $\mathcal{R}$ . It is known (see [34, 29]) that, given the set  $\mathcal{R}(M) \subset C(\partial M)$ , we can endow it, in a constructive way, with a differentiable structure and a metric tensor  $\tilde{g}$ , so that  $(\mathcal{R}(M), \tilde{g})$  becomes a manifold that is isometric to  $(M, g)$ ,

$$(\mathcal{R}(M), \tilde{g}) \cong (M, g).$$

We complete the reconstruction of  $(M, g)$  with the following result:

**Theorem 7.4.** *Let the induced bundle structure (31) on  $\partial M$  and the response operator  $\Lambda_\Gamma$  be given. Then, for any  $h \in C(\partial M)$ , we can find whether  $h \in \mathcal{R}(M)$  or not.*

*Proof:* Consider a function  $h \in C(\partial M)$ . We have that  $h \in \mathcal{R}(M)$  if and only if there is  $x \in M$  such that  $h = r_x$ . Observe that if such  $x$  exists then, for any  $\varepsilon > 0$ , we have  $h(x) - \varepsilon < d(y, z) < h(x) + \varepsilon$  for any  $z \in \partial M$  and  $y \in B(x, \varepsilon)$ , where  $B(x, \varepsilon) \subset M$  is  $x$ -centered ball of radius  $\varepsilon$ . Therefore, for any  $J \in \mathbb{Z}_+$ , any set of points  $z_j \in \partial M$ , and non-negative numbers  $\tau_j^-, \tau_j^+$  satisfying  $\tau_j^- < h(z_j) < \tau_j^+$ ,  $j = 1, \dots, J$ , the set

$$(81) \quad S = S(\{\Sigma_j, \tau_j^-, \tau_j^+\}_{j=1}^J) \subset M \quad \text{contains an open ball,}$$

if  $\Sigma_j \subset \partial M$  are sufficiently small neighborhoods of  $z_j$ .

On the other hand, assume that for all  $J$ ,  $z_j \in \partial M$ ,  $\tau_j^-, \tau_j^+$  satisfying  $\tau_j^- < h(z_j) < \tau_j^+$ , and sufficiently small  $\Sigma_j$ , condition (81) is satisfied. Then there is a point  $x \in M$  such that

$$|d(x, z_j) - h(z_j)| \leq \tau_j^+ - \tau_j^- + \text{diam}(\Sigma_j), \quad j = 1, 2, \dots, J.$$

Thus, letting  $\tau_j^- \rightarrow h(z_j)$ ,  $\tau_j^+ \rightarrow h(z_j)$  and,  $\text{diam}(\Sigma_j) \rightarrow 0$ , we see that there is  $x_J \in M$  such that  $d(x_J, z_j) = h(z_j)$  for all  $j = 1, 2, \dots, J$ .

Denote by  $\{z_j\}_{j=1}^\infty$  a countable dense set on  $\partial M$ , and let  $x_J$ ,  $J \in \mathbb{Z}_+$  be the above defined points corresponding to the first  $J$  points  $z_j$ ,  $j = 1, \dots, J$ , of this set. This gives us a sequence  $(x_J)_{J=1}^\infty$ . As  $M$  is compact, this sequence has a limit point  $x \in M$ . By density of  $\{z_j\}_{j=1}^\infty$ ,  $h = r_x$ .

By Theorem 7.3, condition (81) is equivalent to the condition

$$\dim(Z(\{\Sigma_j, \tau_j^-, \tau_j^+\}_{j=1}^J)) \neq 0.$$

By Theorem 7.2, this can be verified using the boundary data.  $\square$

Combining the above theorem with Lemma 2.5, we obtain the following result:

**Corollary 7.5.** *The induced bundle structure (31) on  $\partial M$  and the Cauchy data set  $C_0(D)$  determine the Riemannian manifold  $(M, g)$  up to an isometry.*

## 8. FOCUSING SEQUENCES. PROOF OF THEOREM 2.9.

**8.1 Waves and delta-distributions.** In the previous section it was shown that, using the induced bundle structure on  $\partial M$  and the response operator  $\Lambda_\Gamma$ , we can control the supports of the waves  $\mathcal{W}\hat{f}(t)$ . In this section the goal is to construct a sequence of sources,  $(\hat{f}_l)_{l=1}^\infty \subset \mathcal{F}^\infty$  such that, when  $l \rightarrow \infty$ , the corresponding waves  $\mathcal{W}\hat{f}_l(t)$  concentrate, at  $t = T_1$ , where  $T_1$  is defined by (76), at a single point  $y \in M^{\text{int}}$ . These waves turn to be useful for reconstructing the bundle  $V$  and the Dirac-type operator  $D$ .

In the following, let  $\underline{\delta}_y$  denote the Dirac delta-distribution at  $y \in M^{\text{int}}$ , i.e.,

$$\int_M \underline{\delta}_y(x) \phi(x) dV_g(x) = \phi(y), \quad \text{for } \phi \in C_0^\infty(M).$$

Since the Riemannian manifold  $(M, g)$  is already found, for any  $y \in M^{\text{int}}$  we can choose a sequence  $\Sigma_{jl} \subset \partial M$ ,  $0 < \tau_{jl}^- < \tau_{jl}^+ \leq T_1 - T$ , so that, for  $S_l = S(\{\Sigma_{jl}, \tau_{jl}^+, \tau_{jl}^-\}_{j=1}^{J(l)})$ ,

$$(82) \quad S_{l+1} \subset S_l, \quad \bigcap_{l=1}^{\infty} S_l = \{y\}.$$

Let  $Z_l = Z(\{\Sigma_{jl}, \tau_{jl}^-, \tau_{jl}^+\}_{j=1}^{J(l)})$  be the corresponding set of the generalized sources defined in Definition 7.1.

**Definition 8.1.** Let  $S_l$ ,  $l = 1, 2, \dots$ , satisfy (82). We call the sequence  $(\hat{f}_l)_{l=1}^\infty$ , with  $\hat{f}_l \in Z_l$ , a *focusing sequence* of generalized sources of the order  $s \in \mathbb{R}_+$  (for brevity, a focusing sequence), if there is a distribution-valued section  $0 \neq A_y \in \mathcal{D}'(M, V)$  supported at the point  $y \in M^{\text{int}}$  such that

$$\lim_{l \rightarrow \infty} \langle \langle \mathcal{W} \hat{f}_l(T_1), \eta \rangle \rangle = \langle \langle A_y, \eta \rangle \rangle, \quad \text{for all } \eta \in \mathcal{D}(|D_\Gamma|^s).$$

**Lemma 8.2.** Assume that the induced bundle structure (31) on  $\partial M$  and the response operator  $\Lambda_\Gamma$  be given. Then, for any  $s \in \mathbb{R}_+$  and any sequence of generalized sources,  $(\hat{f}_l)_{l=1}^\infty \subset \mathcal{F}^\infty$ , we can determine if  $(\hat{f}_l)_{l=1}^\infty$  is a focusing sequence of the order  $s$  or not.

*Proof:* By Theorem 4.3, any  $\eta \in \mathcal{D}(|D_\Gamma|^s)$  can be represented as  $\eta = \mathcal{W} \hat{h}(T_1)$ ,  $\hat{h} \in \mathcal{F}^s$ . Thus, if  $(\hat{f}_l)_{l=1}^\infty$  is a focusing sequence of the order  $s$ , then the limit,

$$(83) \quad \langle \langle A_y, \mathcal{W} \hat{h}(T_1) \rangle \rangle = \lim_{l \rightarrow \infty} \langle \langle \mathcal{W} \hat{f}_l(T_1), \mathcal{W} \hat{h}(T_1) \rangle \rangle,$$

exists for all  $\hat{h} \in \mathcal{F}^s$ . By Theorem 4.1, the existence of this limit can be verified if we are given the induced bundle structure on  $\partial M$  and the operator  $\Lambda_\Gamma$ .

Conversely, assume that we have a sequence  $(\hat{f}_l)_{l=1}^\infty$ , with  $\hat{f}_l \in Z_l$  such that the limit (83) exists for all  $\hat{h} \in \mathcal{F}^s$  and that this limit is non-zero for some  $\hat{h} \in \mathcal{F}^s$ . Then, by the principle of the uniform boundedness, the mappings

$$\eta \mapsto \langle \langle \mathcal{W} \hat{f}_l(T_1), \eta \rangle \rangle, \quad l \in \mathbb{Z}_+,$$

form a uniformly bounded family in the dual of  $\mathcal{D}(|D_\Gamma|^s)$ . By the Banach-Alaoglu theorem, we find a weak\*-convergent subsequence in this space with the limit

$$A_y := \lim_{l \rightarrow \infty} \mathcal{W} \hat{f}_l(T_1) \in \left( \mathcal{D}(|D_\Gamma|^s) \right)'.$$

This gives us the required non-zero distribution-valued section in  $\mathcal{D}'(M, V)$ . As  $\hat{f}_l \in Z_l$ , then  $\text{supp}(A_y) = \{y\}$ .  $\square$

Next we consider the representation of  $A_y$  in a local trivialisation near  $y$ . Since  $\text{supp}(A_y) = \{y\}$ ,  $A_y$  is a finite linear combination, with coefficients from  $\pi^{-1}(y)$ , of the Dirac delta-distribution at  $y$  and its derivatives. The role of the smoothness index  $s$  is just to select the order of this distribution, as is seen in the following result.

**Lemma 8.3.** *Let  $A_y$ ,  $y \in M^{\text{int}}$  be a limit of a focusing sequence of the order  $s$ ,  $\frac{n}{2} < s < \frac{n+1}{2}$ . Then  $A_y$  is of the form*

$$A_y(x) = \lambda \underline{\delta}_y,$$

where  $\lambda \in \pi^{-1}(y)$ . Furthermore, for any  $\lambda \in \pi^{-1}(y)$  there is a focusing sequence  $(\hat{f}_l)_{l=1}^\infty$  such that the waves  $\mathcal{W}\hat{f}_l(T_1) \rightarrow \lambda \underline{\delta}_y$  in  $\mathcal{D}'(M, V)$  when  $l \rightarrow \infty$ .

*Proof:* As for  $s \geq 0$

$$H_{\text{comp}}^s(M, V) \subset \mathcal{D}(|D_\Gamma|^s) \subset H^s(M, V),$$

the embedding theorems imply that, for  $\frac{n}{2} < s < \frac{n+1}{2}$ , the delta-distributions  $\lambda \underline{\delta}_y$ ,  $\lambda \in \pi^{-1}(y)$ , are in  $(\mathcal{D}(|D_\Gamma|^s))'$  while the derivatives of the delta-distribution are not in  $(\mathcal{D}(|D_\Gamma|^s))'$ .

To prove the assertion we have only to verify that a desired focusing sequence exists for any  $\lambda \in \pi^{-1}(y)$ . This can be done by an approximation of  $\lambda \underline{\delta}_y$  by smooth sections with supports converging to  $y$  and an application of the global controllability Theorem 4.3.  $\square$

## 8.2. Construction of the bundle structure.

**Lemma 8.4.** *Assume that we are given the induced bundle structure (31) on  $\partial M$  and the response operator  $\Lambda_\Gamma$ . Then, for any  $y_0 \in M^{\text{int}}$ , we can find integers  $k(1), \dots, k(d)$  and a neighborhood  $U \subset M^{\text{int}}$  of  $y_0$  such that the eigenfunctions  $\phi_{k(1)}(y), \dots, \phi_{k(d)}(y)$  of  $D_\Gamma$  form a basis in  $\pi^{-1}(y)$ ,  $y \in U$ . Moreover, for any  $\hat{h} \in \mathcal{F}^\infty$ , we can find the coefficients  $\alpha_l(y, t)$  of the decomposition*

$$(84) \quad \mathcal{W}\hat{h}(y, t) = \sum_{l=1}^d \alpha_l(y, t) \phi_{k(l)}(y), \quad y \in U, \quad t \in \mathbb{R}.$$

*Proof:* Let  $U_1 \subset M^{\text{int}}$  be a sufficiently small neighborhood of a point  $y_0 \in M^{\text{int}}$  so that there is a trivialisation  $\Phi : \pi^{-1}(U_1) \rightarrow U_1 \times \mathbb{C}^d$ . For all  $y \in U$  we choose  $d$  families of focusing sequences,  $(\hat{f}_k^{j,y})_{k=1}^\infty$ ,  $j = 1, \dots, d$ , such that  $\mathcal{W}\hat{f}_k^{j,y}(T)$  converge to some distributions  $\lambda_y^j \underline{\delta}_y$ ,  $\lambda_y^j \in \pi^{-1}(y)$ ,  $y \in U_1$ .

Using the operator  $\Lambda_\Gamma$  and the induced bundle structure on  $\partial M$  we can evaluate, by Theorem 4.1, the inner products

$$(85) \quad \lim_{k \rightarrow \infty} \langle \langle \mathcal{W}\hat{h}(T), \mathcal{W}\hat{f}_k^{j,y}(T) \rangle \rangle = \langle \langle \mathcal{W}\hat{h}(T), \lambda_y^j \underline{\delta}_y \rangle \rangle = \langle \mathcal{W}\hat{h}(y, T), \lambda_y^j \rangle_y,$$

where  $j = 1, \dots, d$ . The set  $\{\mathcal{W}\hat{h}(T) : \hat{h} \in \mathcal{F}^\infty\}$  contains  $C_0^\infty(M, V)$ . Therefore,  $\{\mathcal{W}\hat{h}(y, T) : \hat{h} \in \mathcal{F}^\infty\} = \pi^{-1}(y)$  and it is possible to verify, using (85), whether the set  $\{\lambda_y^j, j = 1, \dots, d\}$  is a basis in  $\pi^{-1}(y)$  for  $y$  lying in some neighborhood  $U \subset U_1$  of  $y_0$ . Moreover, we can verify whether  $\lambda_y^j$ , considered as functions of  $y$ , define  $C^\infty$ -smooth sections in  $U$ .

On the other hand, Lemma 8.3 guarantees the existence of focusing sequences  $(\hat{f}_l^{j,y})_{l=1}^\infty$  such that the corresponding  $\lambda_y^j$  are linearly independent and smooth in  $U$ .

For any  $\hat{h} \in \mathcal{F}^\infty$ ,  $t \in \mathbb{R}$ , consider the map  $K_t : \mathcal{F}^\infty \rightarrow C^\infty(U, \mathbb{C}^d)$ ,

$$K_t \hat{h}(y) = (\langle \mathcal{W}\hat{h}(y, t), \lambda_y^1 \rangle_y, \dots, \langle \mathcal{W}\hat{h}(y, t), \lambda_y^d \rangle_y), \quad y \in U.$$

By Proposition 5.2, we can find the generalized sources  $\hat{h}_k$ ,  $k = 1, 2, \dots$  such that  $\mathcal{W}\hat{h}_k(T) = \phi_k$ . Thus, for any  $d$  numbers  $k(1), \dots, k(d) \in \mathbb{Z}_+$ , we can evaluate the matrix

$$\mathcal{E}(y) := [e_{j,l}(y)]_{j,l=1}^d, \quad e_{j,l}(y) = \langle \phi_{k(l)}(y), \lambda_y^j \rangle_y, \quad y \in U_1.$$

As for any  $y_0 \in M^{\text{int}}$  the vectors  $\{\phi_k(y_0)\}_{k=1}^\infty$  span  $\pi^{-1}(y_0)$ , it is possible to select  $k(1), \dots, k(d)$  so that the system  $\Phi$  of  $d$  eigenfunctions,  $\Phi = (\phi_{k(1)}, \dots, \phi_{k(d)})$  is linearly independent at  $y_0$ . As  $\mathcal{E}(y)$  is smooth, we can choose  $U \subset U_1$  to be a neighborhood of  $y_0$  where  $\mathcal{E}(y)$  is invertible.

Let  $\hat{h} \in \mathcal{F}^\infty$ . Then the coefficients  $\alpha_i(y, t)$  in (84) can be found by computing the vector  $(\alpha_1(y, t), \dots, \alpha_d(y, t)) = \mathcal{E}(y)^{-1} K_t \hat{h}(y)$ .  $\square$

Next we consider an eigenfunction basis near the boundary. Due to the boundary condition  $P_\Gamma(\phi_k|_{\partial M}) = 0$ , the eigenfunctions  $\phi_k$  do not span  $\pi^{-1}(y_0)$ ,  $y_0 \in \partial M$ . Because of this, we will use near  $\partial M$  a combination of the eigenfunctions  $\phi_k^+$  of  $D_{\Gamma,+}$  and  $\phi_j^-$  of  $D_{\Gamma,-}$ , where  $D_{\Gamma,+} = D_\Gamma$  and  $D_{\Gamma,-} = D_P$  with  $P = (I - P_\Gamma)$  (compare with [2] where a coordinate system near  $\partial M$  was constructed using a combination of the Dirichlet and Neumann eigenfunctions). We denote the spaces of the generalized sources corresponding to the operators  $D_{\Gamma,\pm}$  by  $\mathcal{F}_\pm$  and  $\mathcal{F}_\pm^s$  and by  $\mathcal{W}_\pm$  be propagators (66) for  $D_{P,\pm}$ . We note that our previous results concerning the unique continuation, controllability, and the construction of focusing sequences can be proven using the boundary condition  $I - P_\Gamma = \frac{1}{2}(I - \Gamma)$  instead of  $P_\Gamma = \frac{1}{2}(I + \Gamma)$ .

To simultaneously consider the operators  $D_{\Gamma,+}$  and  $D_{\Gamma,-}$ , we need the following modification of Theorem 4.1:

**Lemma 8.5.** *Assume that we are given the induced bundle structure on  $\partial M$  and the response operator  $\Lambda_\Gamma$ . Then for  $f_+ \in C_0^\infty(\partial M \times \mathbb{R}_+, P_\Gamma W)$  and  $f_- \in C_0^\infty(\partial M \times \mathbb{R}_+, (1 - P_\Gamma)W)$  we can evaluate the inner products*

$$\langle \langle u^{f_+}(t), u^{f_-}(s) \rangle \rangle, \quad \text{for } t, s \geq 0.$$

Here  $u^{f_+}$  is the solution of (33)-(34) with  $P = P_\Gamma$  and  $u^{f_-}$  is the solution of (33)-(34) with  $P = I - P_\Gamma$ .

*Proof:* Let  $I(s, t) = \langle \langle u^{f+}(t), u^{f-}(s) \rangle \rangle$ . Similarly to the proof of Theorem 4.1, we use integration by parts to see that

$$(86) \quad (\partial_s + \partial_t)I(s, t) = i \langle \langle Du^{f+}(t), u^{f-}(s) \rangle \rangle - i \langle \langle u^{f+}(t), Du^{f-}(s) \rangle \rangle \\ = i \int_{\partial M} \langle \gamma(N) \Lambda_{\Gamma}^+ f_+(t), \Lambda_{\Gamma}^- f_-(s) \rangle dA_g.$$

Here  $\Lambda_{\Gamma}^{\pm}$  are the response operators for  $D_{\Gamma, \pm}$ . By Lemma 2.5 the operator  $\Lambda_{\Gamma} = \Lambda_{\Gamma}^+$  and the induced bundle structure determine  $\Lambda_{\Gamma}^-$ . As  $I(0, t) = I(s, 0) = 0$ , we can determine  $I(s, t)$  by solving the differential equation (86) along the characteristics  $t - s = c$ .  $\square$

Let us denote by  $\psi_k$  of the eigenfunctions of  $D_{\Gamma, +}$  and  $D_{\Gamma, -}$ , renumbered so that  $\psi_{2k-1} = \phi_k^+$  and  $\psi_{2k} = \phi_k^-$ ,  $k \geq 1$ . To generalize Lemma 8.4, we need the following result:

**Lemma 8.6.**  $\mathcal{D}(D_{\Gamma, +}^{\infty}) + \mathcal{D}(D_{\Gamma, -}^{\infty}) = C^{\infty}(M, V)$ .

*Proof:* For  $z \in \partial M$  let be a sufficiently small ball  $B = B(z, r) \subset M$  such that the boundary normal coordinates and a local trivialization  $\Phi_B : \pi^{-1}(B) \rightarrow B \times \mathbb{C}^d$  are well defined in  $B$ . It suffices to show that any  $u \in C^{\infty}(M, V)$  with  $\text{supp}(u) \subset B(z, r/2)$  can be written as

$$(87) \quad u = u_+ + u_-, \quad \text{where } u_+ \in \mathcal{D}(D_{\Gamma, +}^{\infty}), \quad u_- \in \mathcal{D}(D_{\Gamma, -}^{\infty}).$$

To prove (87), let  $u^j \in C^{\infty}(\partial M, W)$ ,  $j = 0, 1, 2, \dots$  be supported on  $B(z, r/2) \cap \partial M$ . By using Seeley's extension theorem [51] in  $B$ , there is  $u \in C^{\infty}(M, V)$  supported on  $B(z, r/2)$  such that

$$\partial_N^j u|_{\partial M} = u^j.$$

Moreover, using representations (7), (16) of the Dirac operator in the local trivialization  $\Phi_B$ , we see that

$$(88) \quad D^j u|_{\partial M} = \gamma(N) u^j + A_j(u^0, u^1, \dots, u^{j-1}),$$

where  $A_j$  is a differential operator. Let  $w^j \in C_0^{\infty}(\partial M, W)$ ,  $j = 0, 1, 2, \dots$ , be supported in  $B(z, r/2) \cap \partial M$ . Solving recurrently equations (88) we see that there is  $u \in C^{\infty}(M, V)$  such that  $D^j u|_{\partial M} = w_j$ .

Let  $v \in C^{\infty}(M, V)$ . Then  $v \in \mathcal{D}(D_{\Gamma, \pm}^{\infty})$  if and only if

$$(89) \quad P_{\Gamma, \pm}(D^j v)|_{\partial M} = 0, \quad j = 0, 1, 2, \dots$$

Thus, if  $u \in C^{\infty}(M, V)$  supported in  $B(z, r/2)$  is of the form (87), we have by (89) that

$$D^j u_+|_{\partial M} = (1 - P_{\Gamma})(D^j u|_{\partial M}) =: w_j^+, \\ D^j u_-|_{\partial M} = P_{\Gamma}(D^j u|_{\partial M}) =: w_j^-.$$

On other hand, let  $u \in C^{\infty}(M, V)$  and  $u_+, u_- \in C^{\infty}(M, V)$  be functions such that  $D^j u_{\pm}|_{\partial M} = w_j^{\pm}$ . Then  $u_{\pm} \in \mathcal{D}(D_{\Gamma, \pm}^{\infty})$  and (87) is satisfied.  $\square$

**Lemma 8.7.** *Assume that we are given the induced bundle structure (31) on  $\partial M$  and the response operator  $\Lambda_\Gamma$ . Then, for any  $y_0 \in M$ , we can find integers  $k(1), \dots, k(d)$  and a neighborhood  $U \subset M$  of  $y_0$  such that the eigenfunctions  $\psi_{k(1)}(y), \dots, \psi_{k(d)}(y)$  of  $D_{\Gamma,+}$  or  $D_{\Gamma,-}$  form a basis of  $\pi^{-1}(y)$ ,  $y \in U$ . Moreover, for any  $\hat{h} \in \mathcal{F}_\pm^\infty$ , we can find coefficients  $\alpha_l^\pm(y, t)$  such that*

$$(90) \quad \mathcal{W}_\pm \hat{h}_\pm(y, t) = \sum_{l=1}^d \alpha_l^\pm(y, t) \psi_{k(l)}(y), \quad y \in U, \quad t \in \mathbb{R}.$$

*Proof:* First we show that, for any  $y_0 \in \partial M$ , the space  $\pi^{-1}(y_0)$  is spanned by  $\{\phi_k^+(y_0)\}_{k=1}^\infty \cup \{\phi_k^-(y_0)\}_{k=1}^\infty$ . To this end, let  $u \in C^\infty(M, V)$ . By Lemma 8.6 there are  $u_+ \in \mathcal{D}(D_{\Gamma,+}^\infty)$  and  $u_- \in \mathcal{D}(D_{\Gamma,-}^\infty)$  such that  $u = u_+ + u_-$ . Consider the eigenfunction expansions

$$(91) \quad u_\pm = \sum_{k=1}^\infty a_j^\pm \phi_k^\pm$$

that converge in  $\mathcal{D}(D_{\Gamma,\pm}^\infty)$ . By elliptic regularity in Theorem 2.1, the series (91) converges in  $C^\infty(M, V)$ . In particular, any  $\mu \in \pi^{-1}(y_0)$  can be written in the form  $\mu = \sum_{k=1}^\infty a_j^+ \phi_k^+(y_0) + \sum_{k=1}^\infty a_j^- \phi_k^-(y_0)$ .

Now, let  $y_0 \in \partial M$ . Arguing as in the proof of Lemma 8.4 but using Lemma 8.5 instead of Theorem 4.1, we find  $k(1), \dots, k(d) \in \mathbb{Z}_+$  and neighborhoods  $U$  of  $y_0$  such that  $\psi_{k(1)}(y), \dots, \psi_{k(d)}(y)$  form a smooth basis in  $\pi^{-1}(U \cap M^{\text{int}})$ . Moreover, we can find, using the boundary data, the coefficients  $\alpha_l^\pm(y, t)$  in (90) for any  $y \in U \cap M^{\text{int}}$ . Then, by Lemma 8.6,  $\psi_{k(1)}(y), \dots, \psi_{k(d)}(y)$  form a smooth basis in  $\pi^{-1}(U)$  if and only if the coefficients  $\alpha_l^\pm(y, t)$  may be continued smoothly on the whole  $U$  for any  $\hat{h} \in \mathcal{F}_\pm^\infty$ . This can be verified using the boundary data.  $\square$

The system  $\Phi = (\psi_{k(1)}(y), \dots, \psi_{k(d)}(y))$  of eigenfunctions and the neighborhood  $U$  in Lemma 8.7 define a local trivialization of the vector bundle  $V$ , namely the map  $A_{\Phi,U} : \pi^{-1}(U) \rightarrow U \times \mathbb{C}^d$ ,

$$A_{\Phi,U}(v) = (\alpha_j(y))_{j=1}^d, \quad v \in \pi^{-1}(y), \quad \text{when } v = \sum_{l=1}^d \alpha_l(y) \psi_{k(l)}(y).$$

**Proposition 8.8.** *The induced bundle structure (31) on  $\partial M$  and the Cauchy data set  $C_0(D)$  determine a bundle  $\tilde{V}$  on  $M$  that is bundle morphic to  $V$ .*

*Proof:* Recall that by Lemma 2.5 we know  $\Lambda_\Gamma$ . By Lemma 8.7 we can find, for any  $y_0 \in M$ , a neighborhood  $U$  and a system  $\Phi$  of the eigenfunctions that define a local trivialization in  $\pi^{-1}(U)$ . Thus it remains to prove that the boundary data make it possible to evaluate the transition functions

$$T_{\Phi,U,\tilde{\Phi},\tilde{U}} : (U \cap \tilde{U}) \times \mathbb{C}^d \rightarrow (U \cap \tilde{U}) \times \mathbb{C}^d$$



between any two trivializations  $A_{\Phi,U}$  and  $A_{\tilde{\Phi},\tilde{U}}$ .

Let  $\hat{f}_+ \in \mathcal{F}_+^\infty$ ,  $\hat{f}_- \in \mathcal{F}_-^\infty$ , and  $t \in \mathbb{R}$ . By Lemma 8.7, we can find decompositions (90) of  $\mathcal{W}_+\hat{f}_+(t)$  and  $\mathcal{W}_-\hat{f}_-(t)$  with respect to both  $A_{\Phi,U}$  and  $A_{\tilde{\Phi},\tilde{U}}$ . Therefore, we can find the representations of the function

$$w(y) = \mathcal{W}_+\hat{f}_+(y,t) + \mathcal{W}_-\hat{f}_-(y,t), \quad y \in U \cap \tilde{U}$$

in the both trivializations  $A_{\Phi,U}$  and  $A_{\tilde{\Phi},\tilde{U}}$ . According to Lemma 8.6, the function  $w$  runs over  $C^\infty(M,V)$ , when  $\hat{f}_+$  runs over  $\mathcal{F}_+^\infty$  and  $\hat{f}_-$  over  $\mathcal{F}_-^\infty$ . This makes it possible to compare representations

$$w(y) = \sum_{l=1}^d \alpha_l(y) \psi_{k(l)}(y) = \sum_{l=1}^d \tilde{\alpha}_l(y) \psi_{\tilde{k}(l)}(y), \quad y \in U \cap \tilde{U}$$

when  $w$  runs over  $C^\infty(M,V)$  and, therefore, to determine the transition function  $T_{\Phi,U,\tilde{\Phi},\tilde{U}}$ .  $\square$

**8.3. Construction of operators on  $V$ .** We are now in the position to complete the proof of the main Theorem 2.9.

*Proof:* To this far, we have reconstructed  $(M,g)$  upto an isometry and a bundlemorphic copy  $\tilde{V}$  of  $V$ . To reconstruct the Hermitian and the Clifford module structures and the operators  $F$  and  $D$  on  $\tilde{V}$ , it is enough to find their representations in an arbitrary local trivialisation  $A_{\Phi,U}$ . We will first consider the local trivializations  $A_{\Phi,U}$  where  $U \subset M^{\text{int}}$  and  $\Phi$  consists of eigenfunctions of  $D_{\Gamma,+}$  only. In this connection we will skip index  $+$  in the following considerations.

We start with the reconstruction of the Hermitian structure. Let  $\hat{f} \in \mathcal{F}^\infty$ ,  $y_0 \in U$  and consider a ball  $B(y_0,r) \subset U$ . Denote by  $\mathcal{K}_r = \mathcal{K}_r(\hat{f})$  the set of those  $\hat{h} \in \mathcal{F}^\infty$  for which

$$(92) \quad \langle \langle \mathcal{W}\hat{f}(T) - \mathcal{W}\hat{h}(T), \lambda_y \delta_y \rangle \rangle = 0, \text{ for all } \lambda_y \in \pi^{-1}(y), y \in B(y_0,r).$$

Note that condition (92) is equivalent to the fact that  $\mathcal{W}\hat{h}(T)|_{B(y_0,r)} = \mathcal{W}\hat{f}(T)|_{B(y_0,r)}$ .

Moreover, it follows from Lemma 8.3 that condition (92) can be directly verified from the boundary data. Therefore, using the boundary data we can evaluate

$$(93) \quad \|\mathcal{W}\hat{f}(T)\|_{L^2(B(y_0,r),V)}^2 = \inf_{\hat{h} \in \mathcal{K}_r} \|\hat{h}\|_{\mathcal{F}}^2.$$

As the manifold  $(M,g)$  is already reconstructed and  $\mathcal{W}\hat{f}(T) \in C^\infty(M,V)$ , the equation (93) makes it possible to find

$$(94) \quad |\mathcal{W}\hat{f}(y_0,T)|_{y_0}^2 = \lim_{r \rightarrow 0} \frac{\|\mathcal{W}\hat{f}(T)\|_{L^2(B(y_0,r),V)}^2}{\text{vol}(B(y_0,r))}$$

for an arbitrary  $y_0 \in U \subset M^{\text{int}}$  and  $\hat{f} \in \mathcal{F}_+^\infty$ . Again, as  $\mathcal{W}\hat{f}(T)$  runs over  $\mathcal{D}((D_{\Gamma,+})^\infty)$  when  $\hat{f}$  runs over  $\mathcal{F}_+^\infty$ , it follows from Lemma 8.4

that we can find the Hermitian structure on  $\pi^{-1}(y_0)$  in the trivialization  $A_{\Phi,U}$ . As  $y_0 \in U$  is arbitrary, we can determine the Hermitian structure on  $\tilde{V}|_{M^{\text{int}}}$ .

Next we consider projectors  $\Pi_+ : \pi^{-1}(y_0) \rightarrow \pi^{-1}(y_0)$ ,  $y_0 \in M^{\text{int}}$ . Let  $\hat{f} \in \mathcal{F}^\infty$ . By Theorem 4.1 and Lemma 8.4 we first find, using the boundary data,  $\hat{h} \in \mathcal{F}$  such that  $\Pi_+ \mathcal{W}\hat{f}(T) = \mathcal{W}\hat{h}(T)$ . As  $\mathcal{W}\hat{f}(y_0, T)$  runs over  $\pi^{-1}(y_0)$  when  $\hat{f}$  runs over  $\mathcal{F}$ , this determines  $\Pi_+$  in the trivialization  $A_{\Phi,U}$ . Since  $F = 2\Pi_+ - I$ , the chirality operator  $F$  is determined.

When  $U \subset M^{\text{int}}$  is small enough, there is an orthonormal frame  $(e_1(y), \dots, e_n(y))$  on  $TM|_U$ . Let  $y_0 \in U$  and let  $x(y) = (x^1(y), \dots, x^n(y))$ ,  $x(y_0) = 0$ , be Riemannian normal coordinates centered in  $y_0$  with  $e_l = \partial_l$  at  $y_0$ . By Theorem 4.3 there are generalized sources  $\hat{h}_{jl} \in \mathcal{F}^\infty$ ,  $j = 1, \dots, d$ ,  $l = 1, \dots, n$ , such that

$$u_{jl}(y, T) := \mathcal{W}\hat{h}_{jl}(y, T) = x^l(y)\chi(y)\phi_{k(j)}(y), \quad y \in U.$$

Here  $\chi \in C_0^\infty(U)$  is a cut-off function that is equal to 1 in a neighborhood  $\tilde{U} \subset U$  of  $y_0$ . Moreover, by Lemma 8.4, it is possible to find  $\hat{h}_{jl}$  from the boundary data.

Representing the Dirac operator in this local trivialization (cf. (7), (16)),

$$Du_{jl}(y, T) - \lambda_{k(j)} x^l(y) \phi_{k(j)}(y) = \sum_{p=1}^d \frac{\partial}{\partial x^p} (x^l) \gamma(e_p) \phi_{k(j)}(y), \quad y \in \tilde{U}.$$

In particular, at  $y = y_0$

$$(95) \quad Du_{jl}(y_0, T) = \gamma(e_l) \phi_{k(j)}(y_0).$$

The left-hand side of (95) coincides with  $-i\mathcal{W}(\partial_t \hat{h}_{jl})(y_0, T)$ . By Lemma 8.4 the boundary data determines its representation in the local trivialization  $A_{\Phi,U}$ . Thus we can find  $\alpha_{jl}^p(y_0) \in \mathbb{C}$  such that

$$\gamma(e_l) \phi_{k(j)}(y_0) = \sum_{p=1}^d \alpha_{jl}^p(y_0) \phi_{k(p)}(y_0).$$

This provides the representation of  $\gamma(e_l)$  in the local trivialization  $A_{\Phi,U}$ . Thus we recover the Clifford module structure in  $\tilde{V}|_{M^{\text{int}}}$ .

Next, we consider the representation of the Dirac-type operator  $D$ . For any  $\hat{f} \in \mathcal{F}^\infty$  we can find the representations of  $\mathcal{W}\hat{f}(y, T)$  and  $D\mathcal{W}\hat{f}(y, T) = -i\mathcal{W}(\partial_t \hat{f})(T)$  in a local trivialization  $A_{\Phi,U}$ ,  $U \subset M^{\text{int}}$ . Recall that any functions  $w \in C_0^\infty(U, \tilde{V})$  can be written as  $w = \mathcal{W}\hat{f}(T)$  with some  $\hat{f} \in \mathcal{F}^\infty$ . Therefore, the set of all pairs

$$\{(A_{\Phi,U}(\mathcal{W}\hat{f}(T)|_U), A_{\Phi,U}(-i\mathcal{W}(\partial_t \hat{f})(T)|_U)) : \hat{f} \in \mathcal{F}^\infty\}$$

give the graph of the operator  $D : C_0^\infty(U, \tilde{V}) \rightarrow C^\infty(U, \tilde{V})$  in this local trivialization. This determines  $D$  on  $M^{\text{int}}$ .

To recover the Hermitian and the Clifford module structures and also the operators  $F$  and  $D$  near  $\partial M$ , we recall that we have already found the bundle structure of  $\tilde{V}$  everywhere on  $M$ , i.e. the transition functions between local trivializations. Consider a local trivialization  $A_{U,\Phi}$  where  $U \cap \partial M \neq \emptyset$  and  $\Phi = (\psi_{k(1)}(y), \dots, \psi_{k(d)}(y))$ . By the above considerations, we can find the representation of the Hermitian and the Clifford module structures and the operators  $F$  and  $D$  on  $U \cap M^{\text{int}}$  in this trivialization. However, these representations are smooth up to  $\partial M \cap U$ , and we can continue them to  $\partial M \cap U$ . Thus, we have determined the Hermitian and the Clifford module structures and the operators  $F$  and  $D$  everywhere on  $\tilde{V}$ .  $\square$

## 9. INVERSE PROBLEM SPECTRAL PROBLEM.

**9.1 Time harmonic response operator.** Let  $\{\lambda_k^P, \phi_k^P|_{\partial M}\}_{k=1}^\infty$  be the boundary spectral data corresponding to the self-adjoint Dirac-type operator (3),

$$D_P u = Du, \quad \mathcal{D}(D_P) = \{u \in H^1(M, V) : P(u|_{\partial M}) = 0\},$$

where  $P$  is a (possibly non-local) boundary condition satisfying conditions (19) and (18). Clearly,  $\{\phi_k^P|_{\partial M}\}_{k=1}^\infty$  span the subspace  $(I - P)H^{1/2}(\partial M, W)$  and, therefore, determine  $P$ .

We intend to reconstruct, using the boundary spectral data for  $D_P$ , the fixed-frequency Cauchy data maps,  $\Lambda_\lambda^P$ ,

$$\Lambda_\lambda^P : PL^2(\partial M, W) \rightarrow L^2(\partial M, W), \quad \Lambda_\lambda^P f = u_\lambda^f|_{\partial M}.$$

Here  $u_\lambda^f$  is the solution to the boundary value problem,

$$Du_\lambda^f = \lambda u_\lambda^f, \quad P(u_\lambda^f|_{\partial M}) = f, \quad \lambda \notin \sigma(D_P),$$

where  $\sigma(D_P)$  is the spectrum of  $D_P$ .  $\Lambda_\lambda^P$  is an integral operator with its Schwartz kernel  $\Lambda_\lambda^P(x, y)$  formally defined as an expansion,

$$\Lambda_\lambda^P(x, y) = \sum_{j=1}^\infty \frac{\phi_j^P(x) \overline{\phi_j^P(y)}}{\lambda - \lambda_j}.$$

However, cf. [45, 30] for the scalar Schrödinger operator, this series does not converge so that we should regularize it. We use the same method as was used in [30] for the Laplace operator, namely, we represent the operator  $\Lambda_\lambda^P$  in the form,

$$\Lambda_\lambda^P f = \lim_{\mu \rightarrow +\infty} \Lambda_{i\mu}^P f + \int_{i\mu}^\lambda \partial_\nu (\Lambda_\nu^P f) d\nu.$$

Observe that in the strong operator convergence in  $H^{1/2}(\partial M, W)$ ,

$$\partial_\lambda \Lambda_\lambda^P = \lim_{A \rightarrow +\infty} \partial_\lambda \Lambda_{\lambda, A}^P,$$

where the Schwartz kernel of  $\partial_\lambda \Lambda_{\lambda,A}^P$  is given by

$$\partial_\lambda \Lambda_{\lambda,A}^P(x, y) = - \sum_{|\lambda_j^P| \leq A} \frac{\phi_j^P(x) \overline{\phi_j^P(y)}}{(\lambda - \lambda_j)^2}.$$

Thus it remains to find  $\lim_{\mu \rightarrow +\infty} \Lambda_{i\mu}^P$ .

**Lemma 9.1.** *Let  $D_P$  be an operator of form (3) where  $P$  satisfies conditions (18), (19). Then,*

$$(96) \quad \lim_{\mu \rightarrow +\infty} \Lambda_{i\mu}^P f = f + i\gamma(N)f.$$

*Proof:* Consider the boundary value problem

$$(97) \quad D u_{i\mu}^f = i\mu u_{i\mu}^f, \quad P(u_{i\mu}^f|_{\partial M}) = f \in PC^\infty(\partial M, W).$$

We introduce the boundary geodesic coordinates  $x(y) = (x'(y), x_n(y))$  where  $x_n = \text{dist}(y, \partial M)$  and  $x'$  are the coordinates of the nearest boundary point. Using a special trivialisation of  $V$  near  $\partial M$ , see Appendix, formulae (108) and (109), the Dirac-type operator  $D$  takes the form

$$D = \gamma(N) (\partial_n + A(x', x_n, D')),$$

$$A(x', x_n, D') = -\gamma(N) \sum_{\alpha=1}^{n-1} \gamma(e_\alpha) \nabla_\alpha + \frac{n-1}{2} H(x', x_n) - \gamma(N) Q(x', x_n).$$

Here  $H(x', \tau)$  is the mean curvature of the surface  $\partial M_\tau = \{y \in M : \text{dist}(y, \partial M) = \tau\}$  at the point  $y$  with coordinates  $(x', \tau)$ .

We look for the solution of (97) near  $\partial M$ , using a formal WKB expansion,

$$(98) \quad u_{i\mu}^f(x', x_n) \sim e^{-\mu x_n} \sum_{k=0}^{\infty} \frac{1}{\mu^k} u_k(x', x_n).$$

Substitution of (98) to (97) gives rise to a recurrent system of equations with respect to the powers of  $\mu$ ,

$$(99) \quad -(i + \gamma(N)) u_k + A(x', x_n, D') u_{k-1} + \gamma(N) \partial_n u_{k-1} = 0,$$

where  $u_{-1} = 0$ . This system has to be looked at together with the boundary conditions,

$$(100) \quad P(u_k|_{x_n=0}) = \begin{cases} f, & \text{for } k = 0, \\ 0, & \text{for } k \neq 0. \end{cases}$$

Equation (99) tells us that, for any  $x$ ,  $u_0$  is the eigenspace of  $\gamma(N)$  corresponding to the eigenvalue  $-i$ . We denote the eigenprojectors of  $\gamma(N)$  corresponding to the eigenvalues  $-i$  and  $+i$  by  $p_\gamma^-$  and  $p_\gamma^+$ ,

$$p_\gamma^- = \frac{1}{2}(1 + i\gamma(N)), \quad p_\gamma^+ = \frac{1}{2}(1 - i\gamma(N)).$$

Applying projector  $P$  to (99) with  $k = 0$  and using equation  $\gamma(N)P = (I - P)\gamma(N)$ , we get

$$iPu_0 + \gamma(N)(I - P)u_0 = 0.$$

Letting  $x_n = 0$  in the above equation and using (100), we obtain equation (96). To justify it, we, however, need to consider further terms in the WKB-expansion (98).

If we apply the projector  $p_\gamma^-$  to (99) with  $k = 1$ , we see that

$$(101) \quad -i\partial_n p_\gamma^- u_0 + p_\gamma^- A(x', x_n, D') p_\gamma^- u_0 = 0,$$

where we use that  $\partial_n \gamma(N) = 0$  and  $u_0 = p_\gamma^- u_0$ . Since  $\gamma(e_\alpha) \gamma(N) + \gamma(N) \gamma(e_\alpha) = 0$ , the differential operator  $p_\gamma^- A(x', x_n, D') p_\gamma^-$  is of the order 0. Therefore, (101) is an ordinary differential equation for  $p_\gamma^- u_0$  along the normal geodesic with the initial condition given by (96). This determines  $u_0$ . Next we apply  $p_\gamma^+$  to (99) with  $k = 1$  to obtain  $p_\gamma^+ u_1$ ,

$$2ip_\gamma^+ u_1 = p_\gamma^+ A(x', x_n, D') u_0.$$

Together with (100) with  $k = 1$ , this determines  $u_1|_{x_n=0}$ . Further considerations follow the trend above: applying  $p_\gamma^-$  to (99) with  $k = 2$ , we obtain an ordinary differential equation along the normal geodesic for  $p_\gamma^- u_1$ ; applying  $p_\gamma^+$  to (99) with  $k = 2$  we determine  $p_\gamma^+ u_2$  and, by (100) with  $k = 2$ , the value  $u_2|_{x_n=0}$ , etc.

Take a finite sum  $\exp(-\mu x_n) \sum_{k=0}^l \mu^{-k} u_k$  in (98) and cut it by multiplying with a smooth function  $\chi(x_n)$  equal to 1 near  $\partial M$ . Since  $P(\sum_{k=0}^l \mu^{-k} u_k|_{\partial M}) = f$ , and  $\text{dist}(i\mu, \sigma(D_P)) \geq |\mu|$ , we see that

$$\|u_{i\mu}^f - e^{-\mu x_n} \sum_{k=0}^l \mu^{-k} u_k \chi\|_{H^1(M)} \leq c_l \mu^{-l-1}.$$

implying the desired result on  $\partial M$ .  $\square$

**9.2. From time harmonic data to time domain data.** Having  $\Lambda_\lambda^P$  in our disposal, we reconstruct the response operator  $\Lambda_P$ , see (35). To this end we observe that the set

$$\{u^f|_{\partial M \times \mathbb{R}_+} : u^f \text{ solves (33) with some } f \in PC^\infty(\partial M \times \mathbb{R}_+, W)\},$$

coincides with the Cauchy data set  $C_0(D)$ . Due to causality, to reconstruct  $\Lambda_P$  it is sufficient to find its values only on  $PC_0^\infty(\partial M \times \mathbb{R}_+, W)$ .

Take  $f \in PC_0^\infty(\partial M \times \mathbb{R}_+, W)$ , and let  $T^f \in \mathbb{R}_+$  be such that  $f = 0$  for  $t > T^f$ . Therefore,  $u^f(t) \in \mathcal{D}(D_P)$  for  $t > T^f$ . Moreover, the graph norm of the wave  $u^f(t)$ ,

$$\|u^f(t)\|_{\mathcal{D}(D)}^2 = \|Du^f(t)\|_{L^2(M,V)}^2 + \|u^f(t)\|_{L^2(M,V)}^2,$$

is independent of  $t$ ,  $t > T^f$ . Since this norm is equivalent to the  $H^1(M, V)$ -norm on  $\mathcal{D}(D_P)$ , we see that

$$(102) \quad u^f \in C_{0,b}(\mathbb{R}_+, H^1(M, V)), \quad u^f|_{\partial M \times \mathbb{R}_+} \in C_{0,b}(\mathbb{R}_+, H^{1/2}(\partial M, W)),$$

where  $C_{0,b}(\mathbb{R}_+, X)$  is the class of bounded functions with values in  $X$ , which are equal to 0 near  $t = 0$ .

Consider the Fourier transform,

$$\widehat{u}^f(x, \lambda) = \int_{\mathbb{R}} \exp(-i\lambda t) u^f(x, t) dt.$$

By (102),  $\widehat{u}^f(\cdot, \lambda)$  is an analytic  $H^1(M, V)$ -valued function of  $\lambda$  in the lower half plane,  $\lambda = \nu + i\mu$ ,  $\mu < 0$ . For each  $\lambda$ ,  $\widehat{u}^f(\cdot, \lambda)$  is the solution of the Dirichlet problem (97) with  $\lambda$  instead of  $i\mu$  and  $\widehat{f}(\lambda)$  instead of  $f$  in the boundary condition. From  $\Lambda_\lambda^P$  we find  $\widehat{u}^f(\lambda)|_{\partial M}$ . It follows from (102) that  $\widehat{u}^f(\lambda)|_{\partial M}$  is bounded along the line  $\{\nu + i\mu : \nu \in \mathbb{R}\}$  for any  $\mu < 0$ . Thus,

$$u^f|_{\partial M \times \mathbb{R}_+} = \frac{1}{2\pi} \exp(-\mu t) \int_{\mathbb{R}} \exp(i\nu t) \Lambda_{\nu+i\mu}^P \widehat{f}(\nu + i\mu)|_{\partial M} d\nu.$$

Summarizing, we see that the induced bundle structure on  $\partial M$  and the boundary spectral data  $\{\lambda_k, \phi_k|_{\partial M}\}_{k=1}^\infty$  of a Dirac-type operator  $D_P$ , with  $P$  satisfying (19), (18), determine the Cauchy data set,  $C_0(D)$ . By Theorem 2.9 this implies that the boundary spectral data determine the Riemannian manifold,  $(M, g)$ , up to an isometry, and the vector bundle,  $V$ , with its Hermitian and Clifford module structures, as well as the chirality operator,  $F$  and Dirac-type operator,  $D$ , up to a Dirac bundle morphism. Finally, since  $\phi_k|_{\partial M}$ ,  $k = 1, 2, \dots$ , span the subspace  $PC^\infty(\partial M, W)$ , we can determine the projector  $P$ . Thus we can determine  $D_P$ . This proves Theorem 2.10.  $\square$

## 10. DIRAC-TYPE OPERATOR IN $\mathbb{R}^3$

In this section we apply our results to the classical example of the canonical Dirac-type operator in a bounded domain  $M \subset \mathbb{R}^3$ .

Let  $V = M \times \mathbb{C}^4$  be the trivial bundle over  $M$ , each fiber endowed with the standard Hermitian inner product of  $\mathbb{C}^4$ . The unperturbed Euclidean Dirac operator,  $D_0$  has the form

$$(103) \quad D_0 u = i \sum_{k=1}^3 \alpha_k \partial_k u + mc^2 \alpha_0 u, \quad u \in H^1(\mathbb{R}^3; \mathbb{C}^4).$$

Here  $\alpha_\nu$ ,  $\nu = 0, \dots, 3$ , are the standard Dirac matrices,

$$\alpha_0 = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}, \quad \alpha_k = \begin{pmatrix} 0 & \sigma_k \\ \sigma_k & 0 \end{pmatrix}, \quad k = 1, 2, 3,$$

where  $\sigma_k$  are the Pauli  $2 \times 2$  matrices. Then

$$\alpha_\nu \alpha_\mu + \alpha_\mu \alpha_\nu = \begin{cases} 2I, & \text{for } \nu = \mu, \\ 0, & \text{for } \nu \neq \mu. \end{cases}, \quad \nu, \mu = 0, \dots, 3.$$

and the operator (103) falls into the category of the Dirac-type operators considered in this paper with  $F = \alpha_0 \alpha_1 \alpha_2 \alpha_3$  and  $\gamma(e_k) = i\alpha_k$ .

A perturbation  $Q$  of a physical nature, consisting of a zero components of the Lorentz scalar and vector potentials, see e.g. [47], defines a perturbed Dirac operator  $D_{\mathbf{a},q} = D_0 + Q$ . Namely,

$$(104) \quad D_{\mathbf{a},q}u = i \sum_{k=1}^3 \alpha_k (\partial_k + ia_k(x))u + mc^2 \alpha_0 u + \alpha_0 q(x)u,$$

where  $a_k(x), q(x)$  are  $C^\infty$ -smooth real-valued functions which correspond to the magnetic vector potential  $\mathbf{a}(x) = a_1 dx^1 + a_2 dx^2 + a_3 dx^3 \in \Omega^1(M)$  and the scalar potential  $q(x)$ . Then the potential  $Q$  satisfies conditions (11) and (12) and, applying the previous constructions of the paper, we obtain the following theorem:

**Theorem 10.1.** *Let  $M \subset \mathbb{R}^3$  be a bounded domain with a smooth boundary. Let  $D_{\mathbf{a},q}$  and  $D_{\tilde{\mathbf{a}},\tilde{q}}$  be two Dirac-type operators of form (104) in  $M$ . Assume that the Cauchy data sets of these operators are the same, i.e.*

$$C_0(D_{\mathbf{a},q}) = C_0(D_{\tilde{\mathbf{a}},\tilde{q}}).$$

*Then  $\tilde{q} = q$  and*

$$(105) \quad \tilde{\mathbf{a}} - \mathbf{a} = d\Phi + \sum_{m=1}^{b_1} n_m h_m, \quad n_m \in \mathbb{Z},$$

*where  $\{h_m\}_{m=1}^{b_1}$  is a basis of the relative harmonic 1-forms on  $M$ ,  $b_1$  is the first relative Betti number of  $M$  and  $d\Phi$  is the exterior differential of a function  $\Phi \in C^\infty(M, \mathbb{R})$ ,  $\Phi|_{\partial M} = 0$ .*

The 1-forms  $h_m$  are normalized so that if  $\partial M$  has the external component  $\Sigma_0$  and the internal components  $\Sigma_j$ ,  $j = 1, \dots, b_1$ , then for a path  $\eta = \eta([0, 1]) \subset M$ ,  $\eta(0) \in \Sigma_0$ , and  $\eta(1) \in \Sigma_j$  we have  $\int_\eta h_m = \delta_{jm}$ .

*Proof:* Clearly, the standard Hermitian structure of  $\mathbb{C}^4$ ,  $\gamma(e_k) = i\alpha_k$ , and the canonical differential,  $\nabla_{e_k} = \partial_k$ , define the structure of a Dirac bundle in  $M \times \mathbb{C}^4$  with  $D_{\mathbf{a},q}$  and  $D_{\tilde{\mathbf{a}},\tilde{q}}$  being Dirac-type operators on this bundle.

By Theorem 2.9, there is a bundlemorphism,  $L : M \times \mathbb{C}^4 \rightarrow M \times \mathbb{C}^4$  that is a fiberwise isomorphism and, therefore, is given by a smooth matrix-valued function  $L(x)$  which is invertible for all  $x \in M$ . Applying again Theorem 2.9, we see that  $D = L^{-1}\tilde{D}L$ ,  $F = L^{-1}\tilde{F}L$ , and  $\gamma(e_k) = L^{-1}\gamma(e_k)L$  for  $k = 1, 2, 3$ . These imply that  $\alpha_\nu L = L\alpha_\nu$ ,  $\nu = 0, \dots, 3$ , which means, in turn, that  $L(x)$  is a diagonal matrix. As the bundlemorphism  $L$  preserves the Hermitian structure in each fiber, there is a complex valued function  $\kappa(x)$  such that

$$(106) \quad L(x) = \kappa(x)I, \quad |\kappa(x)| = 1.$$

Since the induced bundle structures on the boundary coincide,  $L(x) = I$  for  $x \in \partial M$ . Together with (106) this yields that, cf. [33],  $\kappa(x) =$

$\exp(i\Psi(x))$ , with  $\Psi \in C^\infty(M, \mathbb{R})$  such that  $\Psi(x) = 0$  on the external component of  $\partial M$  and  $\Psi(x) = 2\pi n_m$ ,  $n_m \in \mathbb{Z}$  on the internal  $m$ -th component of  $\partial M$ . The family of bundlemorphisms (106) are precisely those bundlemorphisms that preserve the bundle structure and, in particular, the structure (104) of the Dirac operator.

Since  $D = L^{-1}\tilde{D}L$ , where  $L$  is of form (106), we see that  $\tilde{q} = q$  and

$$(107) \quad \tilde{\mathbf{a}} = \mathbf{a} + d\Psi.$$

We obtain formula (105) from (107) by writing  $d\Psi = d\Phi + \sum_{m=1}^{b_1} n_m h_m$  where  $\Phi|_{\partial M} = 0$ ,  $b_1$  is the first relative Betti number of  $M$  and  $h_m$  is a basis of the harmonic 1-forms on  $M$  satisfying the relative boundary condition (29).  $\square$

Theorem 10.1 is a improvement, for the considered class of perturbations and domain  $M \subset \mathbb{R}^3$ , of the results in [46]. Note that Theorem 10.1 describes, for the Dirac operator, the relation between the inverse problems and the Aharonov-Bohm effect. (For the previous results on the Aharonov-Bohm effect for a Schrödinger operator, see e.g. [15, 57] for the Euclidean Laplacian and [28] for a more general scalar operators on a Riemannian manifold.) In physical terms, Aharonov-Bohm effect means that not only the the magnetic field  $d\mathbf{a}$ , i.e.  $\text{curl}(\mathbf{a})$ , but also the magnetic vector potential  $\mathbf{a}(x)$  has an effect on the boundary measurements. Theorem 10.1 characterizes what part of  $\mathbf{a}$  besides  $d\mathbf{a}$  can be observed from external measurements.

**Appendix. Local Boundary Condition.** In this appendix we provide a trivialisation of  $V$  near the boundary needed in Sections 2 and 9, and outline the proof that the Dirac-type operator with the local boundary condition is self-adjoint. Here, we consider the unperturbed Dirac operator

$$D_\Gamma u = D_0 u, \quad \mathcal{D}(D_\Gamma) = \{u \in H^1(M, V) : P_\Gamma(u|_{\partial M}) = 0\},$$

where  $D_0$  is defined by (7) and the projector  $P_\Gamma$  by (26), (28).

First we consider the needed trivialisation of  $V$ . In the following, we do not change notations for  $\gamma(v)$ ,  $v \in T_x(M)$ ,  $F$ , etc. when changing from one trivialisation to another.

Let  $U \subset M$  be a coordinate chart near  $\partial M$  with boundary normal coordinates,  $x(y) = (x'(y), x_n(y))$ . We may assume that  $x(U) = B' \times [0, \varepsilon]$ , where  $B' \subset \mathbb{R}^{n-1}$  is open. Let  $w_j(x) \in \pi^{-1}(x)$ ,  $x \in U \cap \partial M$ , be smooth sections such that  $\{w_j(x)\}_{j=1}^d$  is an orthonormal basis of  $\pi^{-1}(x)$ ,  $x \in U \cap \partial M$ . We can continue  $w_j$  along the normal geodesics so that  $\nabla_N w_j = 0$ , and obtain sections  $w_j \in C^\infty(U, V)$  that form an orthonormal basis for  $\pi^{-1}(x)$ ,  $x \in U$ . The map

$$\Psi : \left( x, \sum_{j=1}^d a_j(x) w_j(x) \right) \mapsto (x, a_1(x), \dots, a_d(x))$$



defines a local trivialisation  $\Psi : \pi^{-1}(U) \rightarrow U \times \mathbb{C}^d$ . In this trivialisation  $\nabla_N u$  is represented by  $\partial_n u$ .

Denoting  $\Psi(x, v) = (x, \psi(x, v))$ , we can define another trivialisation  $\Phi(x, v) = (x, \omega(x)\psi(x, v))$ , where in the boundary normal coordinates

$$\omega(x', x_n) = \exp \left( -\frac{n-1}{2} \int_0^{x_n} H(x', \tau) d\tau \right)$$

with  $H(x', \tau)$  being the mean curvature of the hypersurface  $\partial M_\tau = \{x \in M : \text{dist}(x, \partial M) = \tau\}$ . In this trivialisation  $D_0$  has the form

$$(108) \quad D_0 = \gamma(N)[\partial_n + A(x', \partial_{x'}; x_n)],$$

where  $A(x_n) = A(x', \partial_{x'}; x_n)$  is the first-order differential operator with respect to  $x'$  which depends on  $x_n$  as a parameter. Namely,

$$(109) \quad A(x', \partial_{x'}; x_n) = -\gamma(N) \sum_{\alpha=1}^{n-1} \gamma(e_\alpha) \nabla_\alpha + \frac{n-1}{2} H(x', x_n).$$

Here,  $\nabla_\alpha = \nabla_{e_\alpha}$ . The obtained operator  $A(x', \partial_{x'}; 0)$  is called the *hypersurface Dirac operator*.

Next we show that  $D_\Gamma$  fits into the framework of the theory developed in [13] implying its self-adjointness and regularity. (For other proofs of the self-adjointness see [16, 22].) To this end, we use the above trivialisation  $\Phi$  of  $V$  near  $\partial M$  so that  $D_0$  has form (108), (109). We first show that  $A = A(\tau) : L^2(H) \rightarrow L^2(H)$  satisfies

$$(110) \quad \gamma(N)A + A\gamma(N) = 0 \quad \text{and} \quad A(0) = A^*(0).$$

To prove the first identity in (110), we introduce the Riemannian normal coordinates on  $\partial M_\tau$  centered at  $x'_0$  which are associated with the orthonormal frame  $(e_1, \dots, e_{n-1})$ . Then,

$$(111) \quad \gamma(N)A = -\gamma(N) \sum_{\alpha=1}^{n-1} \gamma(N)\gamma(e_\alpha)\nabla_\alpha + \frac{n-1}{2} H(x', x_n)\gamma(N).$$

At the point  $(x'_0, \tau)$ ,

$$\nabla_\alpha(\gamma(N)u) = \gamma(N)\nabla_\alpha u + \gamma(\nabla_\alpha N)u = \gamma(N)\nabla_\alpha u - \sum_{\beta=1}^{n-1} S_{\alpha\beta}\gamma(e_\beta)u,$$

where  $S_{\alpha\beta}$  is the 2nd fundamental form of  $\partial M_\tau$  in the introduced normal coordinates centered at  $x'_0$ . Returning to (111), this implies that

$$\gamma(N)A = \gamma(N) \left( \sum_{\alpha,\beta=1}^{n-1} S_{\alpha\beta}\gamma(e_\alpha)\gamma(e_\beta) + \sum_{\alpha=1}^{n-1} \gamma(e_\alpha)\nabla_\alpha\gamma(N) + \frac{n-1}{2} H \right).$$

By (5),  $\gamma(e_\alpha)\gamma(e_\beta) + \gamma(e_\beta)\gamma(e_\alpha) = -2\delta_{\alpha\beta}$  and  $S_{\alpha\beta}$  is symmetric. Therefore, the above equation and (111) yield the first identity in (110).

Similar arguments, employing that in the boundary normal coordinates,  $\nabla_\alpha e_\beta = S_{\alpha\beta}N$ ,  $N = \partial_n$ , show the second identity in (110).

It remains to show that  $P_\Gamma$  satisfies (18) and (19). As  $P_\Gamma = \frac{1}{2}(I + \Gamma)$ , it follows from (5), (8) that  $P_\Gamma$  satisfies (18). To check (19), let  $P_{APS}$  be a spectral projector of  $A(0)$  satisfying (21). First equation of (110) and (8) imply that  $A(I + \Gamma) = (I - \Gamma)A$ . Thus if  $\lambda_k$  is an eigenvalue of  $A(0)$  with an eigenfunction  $\phi_k$ , then

$$\psi_k = P_\Gamma \phi_k - (I - P_\Gamma) \phi_k$$

is an eigenfunction for the eigenvalue  $-\lambda_k$ . In particular, when  $\lambda_k \neq 0$ ,  $\langle \phi_k, \psi_k \rangle = 0$ , so that  $\|P_\Gamma \phi_k\| = \|(I - P_\Gamma) \phi_k\|$  and  $\|\phi_k\| = \|\psi_k\|$ . Consider the decomposition,  $L^2(\partial M, W) = \mathbb{H}_1 \oplus \mathbb{H}_2$ , where  $\mathbb{H}_2$  is the 0-eigenspace of  $A(0)$ . The above equations imply that  $\mathbb{H}_1, \mathbb{H}_2$  are invariant subspaces for  $P_\Gamma$  and

$$\|P_\Gamma - P_{APS}\| = \frac{1}{\sqrt{2}} \quad \text{on } \mathbb{H}_1.$$

As  $\mathbb{H}_2$  is finite-dimensional, then 1 and  $-1$  are not in the essential spectrum of  $P_\Gamma - P_{APS}$ . By [13], Proposition 3.3,  $(P_\Gamma, P_{APS})$  is a Fredholm pair, so that  $D_\Gamma$  is self-adjoint.

#### REFERENCES

- [1] Atiyah M. F., Patodi V. K., Singer I.M. Spectral asymmetry and Riemannian geometry, I, *Math. Proc. Camb. Philos. Soc.*, **77** (1975), 43-69; II, *ibid*, **78** (1975), 405-432; III, *ibid*, **79** (1976), 71-99.
- [2] Anderson M, Katsuda A, Kurylev Y, Lassas M and Taylor M. Boundary regularity for the Ricci equation, geometric convergence, and Gel'fand inverse boundary problem *Invent. Math.*, **105** (2004), 261-321.
- [3] Astala, K., Päivärinta, L. Calderon's inverse conductivity problem in the plane. *Ann. of Math.*, **163** (2006), 265-299.
- [4] Astala, K., Päivärinta, L., Lassas, M. Calderon's inverse problem for anisotropic conductivity in the plane. *Comm. PDE*, **30** (2005), 207-224.
- [5] Belishev, M. An approach to multidimensional inverse problems for the wave equation. (Russian) *Dokl. Akad. Nauk SSSR*, **297** (1987), 524-527.
- [6] Belishev, M., Kurylev, Y. A nonstationary inverse problem for the multidimensional wave equation "in the large". (Russian) *Zap. Nauchn. Sem. (LOMI)* **165** (1987), 21-30, transl. in *J. Soviet Math.* **50** (1990), 1944-1951
- [7] Belishev, M., Kurylev, Y. To the reconstruction of a Riemannian manifold via its spectral data (BC-method). *Comm. PDE*, **17** (1992), 767-804.
- [8] Berline, N., Getzler, E., Vergne, M. *Heat kernels and Dirac operators*. Springer-Verlag, 1992. viii+369 pp.
- [9] Bismut, J.-M., Cheeger, J. Families index for manifolds with boundary, superconnections, and cones. I. Families of manifolds with boundary and Dirac operators. *J. Funct. Anal.*, **89** (1990), 313-363.
- [10] Blagovestchenskii, A. A one-dimensional inverse boundary value problem for a second order hyperbolic equation. *Zap. Nauchn. Sem. LOMI*, **15** (1969), 85-90.
- [11] Booss-Bavnbek, B., Wojciechowski, K. *Elliptic boundary problems for Dirac operators*. Birkhauser, 1993. xviii+307 pp.
- [12] Brüning, J.  $L^2$ -index theorems on certain complete manifolds. *J. Diff. Geom.*, **32** (1990), 491-532
- [13] Brüning, J, Lesch, M. On boundary value problems for Dirac type operators. I. Regularity and self-adjointness. *J. Funct. Anal.*, **185** (2001), 1-62.

- [14] Daskalov, V., Khristov, E. Explicit formulae for the inverse problem for the regular Dirac operator. *Inverse Problems*, **16** (2000), 247-258
- [15] Eskin, G. Inverse boundary value problems and the Aharonov-Bohm effect. *Inverse Problems*, **19** (2003), 49-62.
- [16] Farinelli, S., Schwarz, G. On the spectrum of the Dirac operator under boundary conditions. *J. Geom. Phys.*, **28** (1998), 67-84.
- [17] Gilbarg D., Trudinger N. *Elliptic partial differential equations of second order*. Springer-Verlag, 1983. xiii+513 pp.
- [18] Gilkey P. Curvature and the eigenvalues of the Laplacian for elliptic complexes. *Adv. Math.*, **10** (1973), 344-382.
- [19] Gilkey P. On the index of geometrical operators for Riemannian manifolds with boundary. *Adv. Math.*, **102** (1993), 129-183.
- [20] Grebert, B. Inverse scattering for the Dirac operator on the real line. *Inverse Problems*, **8** (1992), 787-807.
- [21] Hachem, G. The  $\bar{\partial}$  approach to inverse scattering for Dirac operators. *Inverse Problems*, **11** (1995), 123-146.
- [22] Hijazi, O., Montiel, S., Zhang, X. Eigenvalues of the Dirac operator on manifolds with boundary. *Comm. Math. Phys.*, **221** (2001), 255-265.
- [23] Hijazi O., Montiel S., Roldan A. Eigenvalue boundary problems for the Dirac operator. *Comm. Math. Phys.*, **231** (2002), 375-390.
- [24] Horvath, M. On the inverse spectral theory of Schrödinger and Dirac operators. *Trans. Amer. Math. Soc.*, **353** (2001), 4155-4171.
- [25] Imanuvilov, O. Y., Yamamoto, M. Global uniqueness and stability in determining coefficients of wave equations. *Comm. PDE*, **26** (2001), 1409-1425.
- [26] Isozaki, H. Inverse scattering theory for Dirac operators. *Ann. Inst. H. Poincaré Phys. Theor.*, **66** (1997), 237-270.
- [27] Jung, W. Geometrical approach to inverse scattering for the Dirac equation. *J. Math. Phys.*, **38** (1997), 39-48.
- [28] Kachalov, A., Kurylev, Y. Multidimensional inverse problem with incomplete boundary spectral data. *Comm. PDE*, **23** (1998), 55-95.
- [29] Katchalov, A., Kurylev, Y., Lassas, M. Inverse Boundary Spectral Problems, Chapman Hall/CRC Monographs, 2001, 290 pp.
- [30] Katchalov, A., Kurylev, Y., Lassas, M., Mandache, N. Equivalence of time-domain inverse problems and boundary spectral problems. *Inverse Problems*, **20** (2004), 419-436.
- [31] Kawashita M., Kurylev Y., Soga H. Harmonic moments and an inverse problem for the heat equation. *SIAM J. Math. Anal.*, **32** (2000), 522-537
- [32] Kurylev, Y. Admissible groups of transformations that preserve the boundary spectral data in multidimensional inverse problems. (Russian) *Dokl. Akad. Nauk*, **327** (1992), 322-325, transl. *Soviet Phys. Dokl.* **37** (1993), 544-545.
- [33] Kurylev, Y. An inverse boundary problem for the Schrödinger operator with magnetic field. *J. Math. Phys.*, **36** (1995), 2761-2776.
- [34] Kurylev Y. Multidimensional Gel'fand inverse problem and boundary distance map, *Inverse Problems Related with Geometry* (ed. H.Soga)(1997), 1-15.
- [35] Kurylev, Y., Lassas, M. Gelf'and inverse problem for a quadratic operator pencil. *J. Funct. Anal.*, **176** (2000), 247-263.
- [36] Kurylev, Y., Lassas, M., Somersalo, E. Reconstruction of manifold from electromagnetic boundary measurements, *Contemp. Math.*, **333** (2003), 147-162.
- [37] Kurylev, Y., Lassas, M., Somersalo, E. Focusing waves in electromagnetic inverse problems *Contemp. Math.*, **348** (2004), 11-22.

- [38] Kurylev, Y., Lassas, M., Somersalo, E. Maxwell's equations with a polarization independent wave velocity: Direct and inverse problems, to appear in *J. Math. Pures Appl.*
- [39] Lassas M., Uhlmann G. On determining a Riemannian manifold from the Dirichlet-to-Neumann map. *Ann. Sci. Ecole Norm. Sup.*, **34** (2001), 771-787.
- [40] Lasiecka, I., Triggiani, R. *Control Theory for Partial Differential Equations, vol. II*. Cambridge Univ. Press, 2000, 645-1067.
- [41] Melrose, R. *The Atiyah-Patodi-Singer Index Theorem*, A.Peters, 1993, 377 pp.
- [42] Melrose, R., Piazza, P. Families of Dirac operators, boundaries and the  $b$ -calculus. *J. Differential Geom.* **46** (1997), 99-180.
- [43] Müller, W. On the  $L^2$ -index of Dirac operators on manifolds with corners of codimension two, I, *J. Diff. Geom.*, **44** (1996), 97-177.
- [44] Nachman, A. Reconstructions from boundary measurements. *Ann. of Math.*, **128** (1988), 531-576.
- [45] Nachman, A., Sylvester, J. Uhlmann, G. An  $n$ -dimensional Borg-Levinson theorem. *Comm. Math. Phys.*, **115** (1988), 595-605.
- [46] Nakamura, G., Tsuchida, T. Uniqueness for an inverse boundary value problem for Dirac operators. *Comm. PDE*, **25** (2000), 1327-1369.
- [47] Nogami Y., van Dijk W. Dirac equation with a separable potential. *Phys. Rev. C*, **34** (1986), 1855-1859.
- [48] Novikov, R., Khenkin, G. The  $\bar{\partial}$ -equation in the multidimensional inverse scattering problem. (Russian) *Uspekhi Mat. Nauk* **42** (1987), 93-152.
- [49] Pestov, L., Uhlmann, G. Two dimensional compact simple Riemannian manifolds are boundary distance rigid. *Ann. of Math.*, **161** (2005), 1093-1110.
- [50] Russell D. Controllability and stabilizability theory for linear partial differential equations. *SIAM Rev.* **20** (1978), 639-739.
- [51] Seeley R. T. Extension of  $C^\infty$ -functions defined in a half space. *Proc. Amer. Math. Soc.*, **15** (1964), 625-626.
- [52] Sylvester, J., Uhlmann, G. A global uniqueness theorem for an inverse boundary value problem. *Ann. of Math.*, **125** (1987), 153-169.
- [53] Tataru, D. Unique continuation for solutions to PDEs, between Hörmander's theorem and Holmgren's theorem. *Comm. PDE*, **20** (1995), 855-884.
- [54] Tataru, D. Unique continuation for operators with partially analytic coefficients. *J. Math. Pures Appl.*, **78** (1999), 505-521.
- [55] Taylor, M. *Partial differential equations. II*. Springer-Verlag, 1996. 528 pp.
- [56] Tsuchida, T. An inverse boundary value problem for Dirac operators with small potentials. *Kyushu J. Math.*, **52** (1998), 361-382
- [57] Weder, R. The Aharonov-Bohm effect and time-dependent inverse scattering theory. *Inverse Problems*, **18** (2002), 1041-1056.
- [58] Wilcox, C.H. The domain of dependence inequality for symmetric hyperbolic systems. *Bull. Amer. Math. Soc.*, **70** (1964), 149-154.

YAROSLAV KURYLEV, LOUGHBOROUGH UNIVERSITY, DEPARTMENT OF MATHEMATICAL SCIENCES, LEICESTERSHIRE LE11 3TU, UK

MATTI LASSAS, HELSINKI UNIVERSITY OF TECHNOLOGY, INSTITUTE OF MATHEMATICS, PO BOX 1100, FIN-02015 TKK, FINLAND